

Adult Advanced Life Support

2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations

ABSTRACT: This 2020 *International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations* for advanced life support includes updates on multiple advanced life support topics addressed with 3 different types of reviews. Topics were prioritized on the basis of both recent interest within the resuscitation community and the amount of new evidence available since any previous review. Systematic reviews addressed higher-priority topics, and included double-sequential defibrillation, intravenous versus intraosseous route for drug administration during cardiac arrest, point-of-care echocardiography for intra-arrest prognostication, cardiac arrest caused by pulmonary embolism, postresuscitation oxygenation and ventilation, prophylactic antibiotics after resuscitation, postresuscitation seizure prophylaxis and treatment, and neuroprognostication. New or updated treatment recommendations on these topics are presented. Scoping reviews were conducted for anticipatory charging and monitoring of physiological parameters during cardiopulmonary resuscitation. Topics for which systematic reviews and new Consensuses on Science With Treatment Recommendations were completed since 2015 are also summarized here. All remaining topics reviewed were addressed with evidence updates to identify any new evidence and to help determine which topics should be the highest priority for systematic reviews in the next 1 to 2 years.

Katherine M. Berg, MD
:
Jerry P. Nolan, MBChB
On behalf of the Adult
Advanced Life Support
Collaborators

The full author list is available on page S127.

Key Words: AHA Scientific Statements ■ cardiopulmonary arrest ■ cardiopulmonary resuscitation and emergency cardiac care ■ echocardiography ■ post-cardiac arrest care ■ prognostication ■ sudden cardiac arrest ■ ventricular fibrillation

© 2020 American Heart Association, Inc., European Resuscitation Council, and International Liaison Committee on Resuscitation

<https://www.ahajournals.org/journal/circ>

CONTENTS

Abstract.....	S92
Overview	S93
Topics Reviewed in This 2020 ALS CoSTR	S94
Defibrillation Strategies for Ventricular Fibrillation or Pulseless Ventricular Tachycardia	S95
Anticipatory Defibrillator Charging (ALS 2001: ScopRev).....	S95
Double Sequential Defibrillation (ALS 2003: SysRev)	S96
Automated External Defibrillator Versus Manual Defibrillator (ALS 495: EvUp)	S97
Waveform Analysis for Predicting Successful Defibrillation (ALS 601: EvUp)	S97
Airway, Oxygenation, and Ventilation During CPR	S98
Airway Management During Cardiac Arrest (ALS 576, 783, 432, 496, 711, 714: 2019 SysRev, CoSTR Update)	S98
Confirmation of Correct Tracheal Tube Placement (ALS 469: EvUp)	S98
Oxygen Dose During CPR (ALS 889: EvUp)	S99
Automatic Ventilators Versus Manual Ventilation During CPR (ALS 490: EvUp)	S99
Circulatory Support During CPR	S99
ECPR Versus Manual or Mechanical CPR (ALS 723: 2018 SysRev, 2019 CoSTR)	S99
Physiological Monitoring During CPR	S99
Monitoring Physiological Parameters During CPR (ALS 656: Adopted From Pediatric Task Force ScopRev)	S100
Drugs During CPR, Including Timing of Administration	S100
Vasopressors During Cardiac Arrest (ALS 788, 659, 789, 784, 778: 2019 SysRev, CoSTR).....	S101
Antiarrhythmic Drugs for Cardiac Arrest (ALS 428, 493: 2018 SysRev, CoSTR)	S101
IV Versus IO Drug Delivery (ALS 2046: SysRev)	S102
Steroids During CPR (ALS 433: EvUp)	S103
Buffering Agents for Cardiac Arrest (ALS 483: EvUp)	S103
Drugs for Torsades de Pointes (ALS 457: EvUp)	S103
Intra-arrest Prognostication	S104
Point-of-Care Echocardiography for Prognostication During CPR (ALS 658: SysRev)	S104
ETCO ₂ to Predict Outcome of Cardiac Arrest (ALS 459: EvUp)	S106
Cardiac Arrest in Special Circumstances	S107
Cardiac Arrest Associated With Pulmonary Embolism (ALS 435, 581: SysRev)	S107
Cardiac Arrest in Pregnancy (ALS 436: EvUp)	S108
Opioid Toxicity (ALS 441: EvUp)	S109

Postresuscitation Care	S109
Oxygen Dose After ROSC in Adults (ALS 448: SysRev)	S109
Ventilation Strategy After ROSC in Adults (ALS 571: SysRev)	S111
Postresuscitation Hemodynamic Support (ALS 570: EvUp)	S112
Postresuscitation Steroids (ALS 446: EvUp) ...	S113
Prophylactic Antibiotics After Cardiac Arrest (ALS 2000: SysRev)	S113
Post-Cardiac Arrest Seizure Prophylaxis and Treatment (ALS 431, 868: SysRev)	S114
Targeted Temperature Management (ALS 455, 790, 791, 802, 879: EvUp)	S116
Prognostication in Comatose Patients After Resuscitation From Cardiac Arrest	S117
Clinical Examination for Prognostication (ALS 450, 713, 487: SysRev)	S117
Neurophysiological Tests for Prognostication (ALS 450, 713, 460: SysRev)	S119
Blood Biomarkers for Prognostication (ALS 450, 713, 484: SysRev)	S122
Imaging for Prognostication (ALS 450, 713, 458: SysRev)	S123
ALS CoSTR Topics Not Reviewed in 2020.....	S126
Post-ROSC Percutaneous Coronary Intervention.....	S126
Organ Donation After Cardiac Arrest.....	S126
Manual Defibrillation Topics Not Reviewed in 2020	S126
Circulatory Support Topics Not Reviewed in 2020	S126
Drugs During CPR Topics Not Reviewed in 2020	S127
Intra-arrest Monitoring Topics Not Reviewed in 2020	S127
Special Circumstances Topics Not Reviewed in 2020	S127
Postresuscitation Care Topics Not Reviewed in 2020	S127
Acknowledgments.....	S127
Disclosures.....	S128
References.....	S130

OVERVIEW

The *International Consensus on Cardiopulmonary Resuscitation (CPR) and Emergency Cardiovascular Care (ECC) Science With Treatment Recommendations (CoSTR)* is the fourth in a series of annual International Liaison Committee on Resuscitation (ILCOR) publications. This 2020 CoSTR for advanced life support (ALS) includes new topics addressed by systematic reviews performed within the past 12 months and prioritized by the ALS Task Force. In addition, it includes updates of the ALS treatment recommendations that were

published from 2010 through 2019,^{1–8} as needed, and were based on additional evidence evaluations. As a result, this 2020 CoSTR for ALS is the most comprehensive update since 2010. The 3 major types of evidence evaluation supporting this 2020 publication are the systematic review (SysRev), the scoping review (ScopRev), and the evidence update (EvUp).

The SysRev is a rigorous process following strict methodology to answer a specific question, and each of these ultimately resulted in generation of the task force CoSTR included in this publication. The SysRevs were performed by a Knowledge Synthesis Unit, an Expert Systematic Reviewer, or by the ALS Task Force, and many resulted in separate published SysRevs.

To begin the SysRev, the question to be answered was phrased in terms of the population, intervention, comparator, outcome, study design, time frame (PICOST) format. The methodology used to *identify* the evidence was based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA).⁹ The approach used to *evaluate* the evidence was based on the one proposed by the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) working group.¹⁰ Using this approach, the task force rated as high, moderate, low, or very low the certainty/confidence in the estimates of effect of an intervention or assessment across a body of evidence for each of the predefined outcomes. Randomized controlled trials (RCTs) generally began the analysis as high-certainty evidence, and observational studies generally began the analysis as low-certainty evidence; examination of the evidence by using the GRADE approach could result in downgrading or upgrading of the certainty of evidence. For additional information, refer to “Part 2: Evidence Evaluation Process and Guidelines Development in this supplement.”^{11,11a}

When we have quoted unchanged treatment recommendations from the 2010 CoSTR, the language used differs from that in the GRADE approach because GRADE was not used before 2015.^{12,13}

Draft 2020 CoSTRs for ALS were posted on the ILCOR website¹⁴ for public comment between January 3 and January 4, 2020, with comments accepted through January 18, 2020. These new draft 2020 CoSTR statements for ALS were viewed a total of 4205 times with 11 comments received.

This summary statement contains the final wording of the CoSTR statements as approved by the ILCOR task forces and by the ILCOR member councils after review and consideration of comments posted online in response to the draft 2020 CoSTRs. Within this publication, each topic includes the PICOST as well as the CoSTR, an expanded Justification and Evidence-to-Decision Framework Highlights section, and a list of knowledge gaps requiring future research studies. An evidence-to-decision table is included for each CoSTR

in Appendix A in the Supplemental Materials of this publication.

The second major type of evidence evaluation performed to support this 2020 CoSTR for ALS is a ScopRev, which identifies the extent, range, and nature of evidence on a topic or a question. The ScopRevs were performed by topic experts in consultation with the ALS Task Force. The task force analyzed the identified evidence and determined its value and implications for resuscitation practice or research. The rationale for the ScopRev, the summary of evidence, and task force insights are all highlighted in the body of this publication. The most recent treatment recommendation is included. The task force notes whether the ScopRev identified substantive evidence that may result in a change in ILCOR treatment recommendations. If sufficient evidence was identified, the task force suggested consideration of a future systematic review to supply sufficient detail to support the development of an updated CoSTR. All ScopRevs are included in their entirety in Appendix B in the Supplemental Materials of this publication.

The third type of evidence evaluation supporting this 2020 CoSTR for ALS is an EvUp. EvUps are generally performed for topics previously reviewed by ILCOR to identify new studies published after the most recent ILCOR evidence evaluation, typically through use of search terms and methodologies from previous reviews. These EvUps were performed by task force members, collaborating experts, or by members of council writing groups. The EvUps are cited in the body of this publication with reiteration of the original PICOST (if available) and a note as to whether the evidence suggested the need to consider a SysRev; the existing ILCOR treatment recommendation is quoted. In this publication, no change in ILCOR treatment recommendations resulted from an EvUp; if substantial new evidence was identified, the task force recommended consideration of a SysRev. All EvUps are included in Appendix C in the Supplemental Materials of this publication.

The ALS Task Force considered the availability of new evidence as well as the evidence needed to create, confirm, or revise treatment recommendations. The chapter topics are organized in sections according to the approximate order of the steps of resuscitation, postresuscitation care, and prognostication. For each reviewed topic, the method of review (SysRev, ScopRev, EvUp) is clearly labeled, with links to the relevant review documents in the Appendix.

TOPICS REVIEWED IN THIS 2020 ALS CoSTR

Note: As indicated above, the ALS CoSTR evidence reviews were all completed by January 18, 2020. As a result, this document does not address the topic of

potential influence of coronavirus disease 2019 (COVID-19) on resuscitation practice. In the spring of 2020, an ILCOR writing group was assembled to identify and evaluate the published evidence regarding risks of aerosol generation and infection transmission during attempted resuscitation of adults, children, and infants. This group developed a consensus on science with treatment recommendations and task force insights. This statement is published as a separate document.¹⁵ As new evidence emerges, the ILCOR task forces will review and update this statement, so the reader is referred to the ILCOR website¹⁴ for the most up-to-date recommendations.

Defibrillation Strategies for Ventricular Fibrillation or Pulseless Ventricular Tachycardia

- Anticipatory defibrillator charging (ALS 2001: ScopRev)
- Double sequential defibrillation (ALS 2003: SysRev)
- Automated external defibrillator versus manual defibrillator (ALS 495: EvUp)
- Waveform analysis for predicting successful defibrillation (ALS 601: EvUp)

Airway, Oxygenation, and Ventilation During CPR

- Airway management during cardiac arrest (ALS 576, 783, 432, 496, 711, 714: 2019 SysRev, CoSTR update)
- Confirmation of correct tracheal tube placement (ALS 469: EvUp)
- Oxygen dose during CPR (ALS 889: EvUp)
- Automatic ventilators versus manual ventilation during CPR (ALS 490: EvUp)

Circulatory Support During CPR

- ECPR versus manual or mechanical CPR (ALS 723: 2018 SysRev, 2019 CoSTR)

Physiological Monitoring During CPR

- Monitoring physiological parameters during CPR (ALS 656: Adopted From Pediatric Task Force ScopRev)

Drugs During CPR, Including Timing of Administration

- Vasopressors during cardiac arrest (ALS 788, 659, 789, 784, 778: 2019 SysRev, CoSTR)
- Antiarrhythmic drugs for cardiac arrest (ALS 428, 493: 2018 SysRev, CoSTR)
- Intravenous versus intraosseous drug delivery (ALS 2046: SysRev)
- Steroids during cardiac arrest (ALS 433: EvUp)
- Buffering agents for cardiac arrest (ALS 483: EvUp)
- Drugs for torsades de pointes (ALS 457: EvUp)

Intra-arrest Prognostication

- Point-of-care echocardiography for prognostication during CPR (ALS 658: SysRev)
- ETCO₂ to predict outcome of cardiac arrest (ALS 459: EvUp)

Cardiac Arrest in Special Circumstances

- Cardiac arrest associated with pulmonary embolism (ALS 435, 581: SysRev)
- Cardiac arrest in pregnancy (ALS 436: EvUp)
- Opioid toxicity (ALS 441: EvUp)

Postresuscitation Care

- Oxygen dose after return of spontaneous circulation (ROSC) in adults (ALS 448: SysRev)
- Ventilation strategy after ROSC in adults (ALS 571: SysRev)
- Postresuscitation hemodynamic support (ALS 570: EvUp)
- Postresuscitation steroids (ALS 446: EvUp)
- Prophylactic antibiotics after cardiac arrest (ALS 2000: SysRev)
- Post-cardiac arrest seizure prophylaxis and treatment (ALS 431, 868: SysRev)
- Targeted temperature management (ALS 455, 790, 791, 802, 879: EvUp)

Prognostication in Comatose Patients After Resuscitation From Cardiac Arrest

- Clinical examination for prognostication (ALS 450, 713, 487: SysRev)
- Neurophysiological tests for prognostication (ALS 450, 713, 460: SysRev)
- Blood biomarkers for prognostication (ALS 450, 713, 484: SysRev)
- Imaging for prognostication (ALS 450, 713, 458: SysRev)

DEFIBRILLATION STRATEGIES FOR VENTRICULAR FIBRILLATION OR PULSELESS VENTRICULAR TACHYCARDIA

The task force restricted its review to 2 new topics that were based on trends in current clinical practice. These deal primarily with manual defibrillation in adults. The CoSTRs for the use of automated external defibrillators for adults can be found in Adult Basic Life Support, and for infants and children in Pediatric Life Support.

Anticipatory Defibrillator Charging (ALS 2001: ScopRev)

Rationale for Review

This topic was chosen because the timing of the rhythm check in relation to manual defibrillator charging varies by country and region. The standard method described in the 2010 American Heart Association Guidelines for CPR and ECC¹⁶ and the 2015 European Resuscitation Guidelines¹⁷ consists of briefly pausing compressions to analyze the rhythm then resuming compressions while charging the defibrillator, then pausing compressions

briefly to deliver the shock. With the anticipatory method, the defibrillator is charged near the end of a compression cycle but before the rhythm is checked; then, compressions are paused briefly both to analyze the rhythm and deliver a shock. The ScopRev methodology was chosen given the limited published evidence.¹⁸

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults with cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: Charging the defibrillator before rhythm analysis during manual defibrillation
- Comparator: Charging the defibrillator after rhythm analysis during manual defibrillation
- Outcome: Survival with favorable neurological/functional outcome at discharge, 30 days, 60 days, 180 days, and/or 1 year; survival only at discharge, 30 days, 60 days, 180 days, and/or 1 year; ROSC were defined as critical or important outcomes. Other outcomes were termination of arrhythmia, defibrillation success, preshock pause, postshock pause, perishock pause, hands-off time, hands-on time, compression fraction, inappropriate shocks, shocks during chest compression (shock to rescuer), or any other defibrillation measure.
- Study design: Human and manikin studies were included. RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) were eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) were excluded. In addition, gray literature (evidence not published in traditional journals) was included in this ScopRev.^{19,20}
- Time frame: All years and languages were included. Studies without a title in English were excluded. MEDLINE, Embase, and Cochrane databases were updated to October 7, 2019.

Summary of Evidence

We identified no clinical studies addressing the critical or important outcomes specified in the PICOST question. Three manikin and 1 multicenter retrospective human study were identified. In the only human study,²¹ both methods resulted in relatively short pre- and postshock pauses, whereas anticipatory charging was associated with a shorter total hands-off time in the 30 seconds preceding shock delivery. The results of the 3 manikin studies showed reduced overall pause duration during the compression cycle, but increased pre, post, and perishock pause duration with anticipatory charging.^{22–24}

Task Force Insights

The ScopRev is included in [Supplement Appendix B-1](#). The task force noted that although anticipatory charging can reduce overall chest compression pause duration during the compression cycle, it can increase pre,

post, and perishock pause duration. The clinical relevance of these findings is undetermined. Further high-quality evidence is required to evaluate the relative importance of the different types of pause duration for critical and important patient outcomes, and the role of new defibrillator technologies and methods. There are insufficient data for a SysRev to be of use at this time.

Treatment Recommendation

There was no treatment recommendation on timing of defibrillator charging previously, and in the absence of sufficient evidence, none was added.

Double Sequential Defibrillation (ALS 2003: SysRev)

Rationale for Review

This is a new topic in response to the increasing use of double (dual) sequential defibrillation (DSD). At least 20% of patients with ventricular fibrillation (VF)/pulseless ventricular tachycardia (pVT) will remain in a shockable rhythm after 3 shocks.^{25–28} Survival decreases as the number of defibrillation attempts required increases. DSD, or the use of 2 defibrillators to deliver 2 overlapping shocks or 2 rapid sequential shocks, one with standard pad placement and the other with either anteroposterior or additional anterolateral pad placement, has been suggested as a possible means of increasing VF termination rates.

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults with cardiac arrest in any setting (in-hospital or out-of-hospital) with a shockable rhythm
- Intervention: DSD
- Comparator: Standard defibrillation
- Outcome: Favorable neurological outcome at hospital discharge, survival to hospital discharge or admission, ROSC, or termination of VF
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies with 5 patients or more) are eligible for inclusion.
- Time frame: There was no date restriction, and the literature search was updated to September 27, 2019.
- International Prospective Register of Systematic Reviews (PROSPERO) Registration: CRD42020152575

Consensus on Science

For the critical outcomes of survival with favorable neurological outcome^{29–31} and survival to hospital discharge^{29–34} and the important outcomes of survival to hospital admission,^{29,30,32,33} ROSC,^{29–35} and termination of VF,^{31,34,35} we identified only observational studies. The overall certainty of evidence was rated as very low for all outcomes, primarily because of a very serious risk

of bias. The individual studies were all at a critical or serious risk of bias because of confounding (due to inadequate adjustment for cardiac arrest characteristics and other factors). Because of this and a high degree of heterogeneity, no meta-analyses could be performed, and individual studies were difficult to interpret.³⁶

Treatment Recommendation

We suggest against routine use of a DSD strategy in comparison with a standard defibrillation strategy for cardiac arrest with a shockable rhythm (weak recommendation, very low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-1](#). There is no strong evidence to favor one intervention compared with the other. The evidence available (very low certainty) suggests lower rates of survival and neurological outcome for patients treated with DSD, but any odds ratios (ORs) or other results reported are difficult to interpret given the very high risk of bias.³⁶ There is no consensus standardized approach to double defibrillation, in that a double-dose strategy could be 2 overlapping shocks or 2 sequential shocks. The ALS Task Force discussed whether any potential benefit might arise from increased shock energy, the fact that 2 shocks were delivered sequentially, different pad placement and vector for the second shock, or some other reason. The task force is aware of recently published data from a small pilot RCT comparing standard defibrillation to DSD (adding a second set of defibrillator pads in the anteroposterior position) or to vector change defibrillation (replacing anterolateral pads with anteroposterior pads).³⁷ The study found differences in VF termination (DSD 76%, vector change 82%, and standard placement 66%) and ROSC (DSD 40%, vector change 39%, and standard defibrillation 25%). This pilot RCT was not designed to formally test differences between the groups, and no survival data were reported. These results have informed a larger, ongoing RCT (NCT04080986) that will provide further data about DSD.

Implementation of DSD requires training of staff and availability of defibrillators. It is important to monitor the intervention to determine effectiveness, and to track adverse events such as harm to the patient, defibrillator damage, and the increase in resource utilization.

Knowledge Gap

- High-quality studies comparing DSD with standard defibrillation in terms of survival and neurological outcome at hospital discharge

Automated External Defibrillator Versus Manual Defibrillator (ALS 495: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults who are in cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: Use of an automated external defibrillator or a multifunctional defibrillator in automatic mode
- Comparator: Standard resuscitation (using a manual defibrillator)
- Outcome: Favorable neurological outcome at hospital discharge, survival to hospital discharge or admission, ROSC, or termination of VF
- This topic was last reviewed in 2010.^{43,44} The evidence update is included in [Supplement Appendix C-1](#) and the search conducted was limited to January 2008 to December 2019. We identified 5 observational studies (only 2 of which included a comparison group) and no randomized trials.³⁸⁻⁴² After consideration, a SysRev was not suggested.

Treatment Recommendation

This treatment recommendation (below) is unchanged from 2010.^{43,44}

No significant survival differences have been demonstrated between defibrillation in semiautomatic and manual modes during out-of-hospital or in-hospital resuscitation; however, the semiautomatic mode is preferred because it is easier to use and may deliver fewer inappropriate shocks.

Trained personnel may deliver defibrillation in manual mode. Use of the manual mode enables chest compressions to be continued during charging, thereby minimizing the preshock pause. When using the defibrillator in manual mode, frequent team training and ECG recognition skills are essential.

The defibrillation mode that results in the best outcome will be influenced by the system of care and by provider skills, training, and ECG recognition.

Waveform Analysis for Predicting Successful Defibrillation (ALS 601: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults with cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: Use of techniques for prediction of the likelihood of success of defibrillation (analysis of VF, etc)
- Comparator: Standard resuscitation (without such prediction)
- Outcome: Survival with favorable neurological/functional outcome at discharge, 30 days, 60 days,

180 days, and/or 1 year; survival only at discharge, 30 days, 60 days, 180 days, and/or 1 year; ROSC; termination of VF

- This topic was last reviewed in 2010.^{43,44} Two EvUps were completed for 2020 and are included in [Supplement Appendix C-2a and C-2b](#). The evidence updates restricted the search to January 2008 to January 2020 and identified one large RCT conducted in 2013⁴⁵ and 20 observational studies.^{46–65} In addition, there is an ongoing multicenter RCT of real-time amplitude spectrum area to guide defibrillation (NCT03237910). Although the VF waveform analyses and outcomes studied were highly heterogeneous, given the amount of data available, an updated SysRev was suggested.

Treatment Recommendation

This treatment recommendation (below) is unchanged from 2010.^{43,44}

There is insufficient evidence to support routine use of VF waveform analysis to guide defibrillation management in adult cardiac arrest in- or out-of-hospital.

AIRWAY, OXYGENATION, AND VENTILATION DURING CPR

Airway Management During Cardiac Arrest (ALS 576, 783, 432, 496, 711, 714: 2019 SysRev, CoSTR Update)

Airway management during cardiac arrest was addressed by a 2019 SysRev⁶⁶ and a 2019 CoSTR summary.^{2,3} Consensus on science, justification and evidence-to-decision highlights, and knowledge gaps can be found in the 2019 CoSTR summary.^{2,3}

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults with cardiac arrest from any cause and in any setting (in-hospital or out-of-hospital)
- Intervention: A specific advanced airway management method (eg, tracheal intubation or a supraglottic airway) during cardiac arrest
- Comparator: A different advanced airway management method or no advanced airway management method (eg, bag-mask ventilation) during cardiac arrest
- Outcome: ROSC, survival, or survival with favorable neurological outcome at discharge/28 days or longer
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) that compared at least 2 airway strategies were eligible

for inclusion. Studies with 10 or fewer patients in either group were excluded.

- Time frame: All years and languages were included; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to October 30, 2018.

Treatment Recommendations

We suggest using bag-mask ventilation or an advanced airway strategy during CPR for adults with cardiac arrest in any setting (weak recommendation, low to moderate certainty of evidence).

If an advanced airway is used, we suggest a supraglottic airway for adults with out-of-hospital cardiac arrest (OHCA) in settings with a low tracheal intubation success rate (weak recommendation, low-certainty evidence).

If an advanced airway is used, we suggest a supraglottic airway or tracheal intubation for adults with OHCA in settings with a high tracheal intubation success rate (weak recommendation, very low-certainty evidence).

If an advanced airway is used, we suggest a supraglottic airway or tracheal intubation for adults with in-hospital cardiac arrest (IHCA) (weak recommendation, very low-certainty evidence).^{2,3}

Confirmation of Correct Tracheal Tube Placement (ALS 469: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults with cardiac arrest in any setting (in-hospital or out-of-hospital) requiring tracheal intubation
- Intervention: Use of devices (eg, waveform capnography, CO₂ detection device, esophageal detector device, or tracheal ultrasound)
- Comparator: Not using these devices
- Outcome: Tracheal intubation success
- This topic was last reviewed in 2015.^{1,7} This EvUp is included in [Supplement Appendix C-3](#). An updated SysRev was not considered necessary.

Treatment Recommendations

This treatment recommendation (below) is unchanged from 2015.^{1,7}

We recommend using waveform capnography to confirm and continuously monitor the position of a tracheal tube during CPR in addition to clinical assessment (strong recommendation, low-quality evidence).

We recommend that if waveform capnography is not available, a nonwaveform CO₂ detector, esophageal detector device, or ultrasound in addition to clinical assessment is an alternative (strong recommendation, low-quality evidence).^{1,7}

Oxygen Dose During CPR (ALS 889: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults with cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: Administering a maximal oxygen concentration (eg, 100% by face mask or closed circuit)
- Comparator: No supplemental oxygen (room air) or an alternative supplemental oxygen concentration (eg, 40% to 50%)
- Outcome: Survival with favorable neurological/functional outcome at discharge, 30 days, 60 days, 180 days, and/or 1 year; survival only at discharge, 30 days, 60 days, 180 days, and/or 1 year; ROSC
- This topic was last reviewed in 2015.^{1,7} This EvUp is included in [Supplement Appendix C-4](#) and the search was conducted from October 30, 2013, to December 2, 2019. The search identified 2 observational studies relevant to this topic published since 2015.^{67,68} There are no adult studies of oxygen titration during CPR. An updated SysRev was not considered necessary.

Treatment Recommendation

This treatment recommendation (below) is unchanged from 2015.^{1,7}

We suggest using the highest possible inspired oxygen concentration during CPR (weak recommendation, very low-certainty evidence).

Automatic Ventilators Versus Manual Ventilation During CPR (ALS 490: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults and children in cardiac arrest in any setting (in-hospital or out-of-hospital) and who have advanced airways in place
- Intervention: The use of automatic ventilators
- Comparator: Use of manual ventilation
- Outcome: Ventilation, oxygenation, hands-off time, continuous compressions, survival
- This topic was last reviewed in 2010.^{6,8} An evidence update is included in [Supplement Appendix C-5](#). A search restricted to January 1, 2008, to December 7, 2019, identified 1 very small RCT and 3 observational studies.⁶⁹⁻⁷² An updated SysRev was not considered necessary.

Treatment Recommendation

This treatment recommendation (below) is unchanged from 2010.^{6,8}

There is insufficient evidence to support or refute the use of an automatic transport ventilator over manual

ventilation during resuscitation of the cardiac arrest victim with an advanced airway.

CIRCULATORY SUPPORT DURING CPR

ECPR Versus Manual or Mechanical CPR (ALS 723: 2018 SysRev, 2019 CoSTR)

Extracorporeal CPR (ECPR) was addressed by a 2018 SysRev⁷³ and a 2019 published CoSTR summary.^{2,3} Consensus on Science, Values, Preferences, and Task Force Insights and Knowledge Gaps can be found in the 2019 CoSTR summary.^{2,3}

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults (18 years or older) and children (younger than 18 years) with cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: ECPR, including extracorporeal membrane oxygenation or cardiopulmonary bypass, during cardiac arrest
- Comparator: Manual CPR and/or mechanical CPR
- Outcome: Short-term survival and neurological outcomes (eg, hospital discharge, 28 days, 30 days, and 1 month) and long-term survival and neurological outcomes (eg, 3 months, 6 months, and 1 year)
- Study design: Randomized trials, non-RCTs, and observational studies (cohort studies and case-control studies) with a control group were included. Animal studies, ecological studies, case series, case reports, reviews, abstracts, editorials, comments, and letters to the editor were not included.
- Time frame: All years and languages were included up to May 22, 2018.

Treatment Recommendations

We suggest that ECPR may be considered as a rescue therapy for selected patients with cardiac arrest when conventional CPR is failing in settings in which it can be implemented (weak recommendation, very low-certainty evidence).^{2,3}

PHYSIOLOGICAL MONITORING DURING CPR

The ability to monitor physiological variables and tailor ALS interventions to the patient's precise physiological state is appealing and hence the ongoing interest in this area.

Monitoring Physiological Parameters During CPR (ALS 656: Adopted From Pediatric Task Force ScopRev)

Rationale for Review

Physiological monitoring during CPR, including measurement of end-tidal CO₂ (ETCO₂) and arterial blood pressure among other parameters, is growing in popularity. There is limited evidence to-date on whether use of such parameters improves outcomes. This topic was last updated in 2015.^{1,7} A Pediatric Task Force ScopRev of physiological monitoring during CPR for 2020 also included review of the adult evidence. The adult portion of the ScopRev was included in this update.

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults who are in cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: The use of physiological feedback in regard to CPR quality (eg, arterial catheter, ETCO₂ monitoring, Spo₂ waveforms, or others)
- Comparator: No use of physiological feedback
- Outcome: Survival with favorable neurological/functional outcome at discharge, 30 days, 60 days, 180 days, and/or 1 year; survival only at discharge, 30 days, 60 days, 180 days, and/or 1 year; ROSC
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies). If it is anticipated that there will be insufficient studies from which to draw a conclusion, case series may be included. The minimum number of cases for a case series to be included was set by the task-force at 5. Unpublished studies (eg, conference abstracts, trial protocols) are excluded.
- Time frame: For Step 1, all languages are included if there is an English abstract. We searched articles from 2015 onward. For Step 2, if a SysRev or ScopRev of high quality (as per AMSTAR 2 tool: <https://amstar.ca/Amstar-2.php>) is identified, the search can be limited to beyond data and/or scope of that review.

Summary of Evidence

ETCO₂ or Arterial Blood Pressure Monitoring

The ScopRev is included in [Supplement Appendix B-2a and 2b](#). We identified 1 observational propensity-matched cohort study of adult IHCA by using data from the AHA Get With the Guidelines-Resuscitation registry.⁷⁴ In this study, 3032 physiologically monitored patients (either by ETCO₂ or arterial catheter) were compared with 6064 patients without such monitoring. Those monitored showed a higher rate of ROSC (OR, 1.22 [95% CI, 1.04; 1.43]) but not survival to discharge (OR, 1.04 [95% CI, 0.91; 1.18]) nor survival with

favorable neurological outcome. The study did not specifically look at diastolic blood pressure. Even when an arterial catheter was in place, only about one third reported using the diastolic blood pressure to guide their CPR efforts.

Near-Infrared Spectroscopy

The ScopRev is included in [Supplement Appendix B-2c](#). Two SysRevs were identified; the latest was published in 2018 and comprised studies published before February 2017. The SysRevs concluded that a higher cerebral oxygen saturation measured with near-infrared spectroscopy (NIRS) is associated with a higher chance of ROSC and survival and a lower NIRS is associated with an increased mortality.^{75,76} However, there is no consensus on specific thresholds of cerebral oxygen saturation.⁷⁵ There was a wide overlap of mean or median cerebral oxygen saturation values between patients with and without ROSC, and this was also reflected in the cohort studies.⁷⁷⁻⁷⁹ Only 1 observational study⁸⁰ compared the rates of ROSC with and without NIRS monitoring and found no difference between the groups. All other studies compared NIRS values in patients who achieved ROSC with those without ROSC. Many different NIRS devices with noninterchangeable saturation indices were used across the studies, complicating comparisons.⁸¹ The findings of the observational studies published since February 2017 correlate with those published in both SysRevs.

The ScopRev did not suggest the existence of sufficient new data to proceed to a SysRev.

Task Force Insights

Physiological monitoring during CPR is increasingly popular and potentially useful for both outcome prediction and real-time improvement in CPR quality. The heterogeneity and observational nature of available studies continues to limit the task force's ability to make specific recommendations. The 2015 treatment recommendation is therefore unchanged.^{1,7}

Treatment Recommendation

This treatment recommendation (below) is unchanged from 2015.^{1,7}

We make no treatment recommendation for any particular physiological measure to guide CPR because the available evidence would make any estimate of effect speculative.

DRUGS DURING CPR, INCLUDING TIMING OF ADMINISTRATION

Since the 2015 CoSTR, there have been RCTs of antiarrhythmics and vasopressors during CPR^{82,83} and subsequent publications comparing the intravenous (IV) and intraosseous (IO) route for drugs.^{84,85}

Vasopressors During Cardiac Arrest (ALS 788, 659, 789, 784, 778: 2019 SysRev, 2019 CoSTR)

The topic of vasopressors during cardiac arrest was addressed by a 2019 SysRev⁸⁶ and a published CoSTR summary. Consensus on science, justification and evidence to decision highlights, and knowledge gaps can be found in the 2019 CoSTR summary.^{2,3}

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults (older than 18 years) with cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: Any vasopressor or combination of vasopressors provided intravenously or intraosseously during CPR
- Comparator: No vasopressor, a different vasopressor, or a combination of vasopressors provided intravenously or intraosseously during CPR
- Outcome: Short-term survival (ROSC and survival to hospital admission), midterm survival (survival to hospital discharge, 28 days, 30 days, or 1 month), midterm favorable neurological outcomes (Cerebral Performance Category [CPC] 1–2 or modified Rankin Scale [mRS] score 0–3 at hospital discharge, 28 days, 30 days, or 1 month), and long-term unfavorable and poor (mRS score 4–5) neurological outcomes (after 1 month)
- Study design: Randomized trials, nonrandomized trials, and observational studies (cohort and case-control studies) with a comparison group were included.
- Time frame: All years and languages were included if there was an English abstract to November 23, 2018.

Treatment Recommendations

We recommend administration of epinephrine during CPR (strong recommendation, low to moderate certainty of evidence).

For nonshockable rhythms (pulseless electric activity/asystole), we recommend administration of epinephrine as soon as feasible during CPR (strong recommendation, very low-certainty evidence).

For shockable rhythms (VF/pVT), we suggest administration of epinephrine after initial defibrillation attempts are unsuccessful during CPR (weak recommendation, very low-certainty evidence).

We suggest against the administration of vasopressin in place of epinephrine during CPR (weak recommendation, very low-certainty evidence).

We suggest against the addition of vasopressin to epinephrine during CPR (weak recommendation, low-certainty evidence).^{2,3}

Additional Task Force Commentary

Concerns have been expressed about epinephrine increasing the number of survivors with unfavorable neurological outcome in the PARAMEDIC2 trial (Pre-Hospital Assessment of the Role of Adrenaline: Measuring the Effectiveness of Drug Administration in Cardiac Arrest). The opinion of the ALS Task Force, however, is that any drug that increases the rate of ROSC and survival, but is given after several minutes of cardiac arrest when some degree of neurological damage may already have occurred, will likely increase the number of survivors with both favorable and unfavorable neurological outcome. Determining the likelihood of favorable or unfavorable neurological outcome at the time of starting resuscitation is currently not feasible. Therefore, the task force consensus is that continuing to use a drug that increases survival and focusing efforts on providing earlier CPR, earlier drug administration, and improved postresuscitation care for all patients is likely to increase survival with a favorable neurological outcome.

Antiarrhythmic Drugs for Cardiac Arrest (ALS 428, 493: 2018 SysRev, CoSTR)

This topic was addressed by a 2018 SysRev⁸⁷ and a published 2018 CoSTR summary.^{4,5} Consensus on Science, Values and Preferences, Task Force Insights, and Knowledge Gaps can be found in the 2018 CoSTR summary.^{4,5}

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults and children in cardiac arrest in any setting (in-hospital or out-of-hospital) and a shockable rhythm at any time during CPR or immediately after ROSC
- Intervention: Administration (intravenously or intraosseously) of an antiarrhythmic drug during CPR or immediately (within 1 hour) after ROSC
- Comparator: Administration of another antiarrhythmic drug or placebo or no drug during CPR or immediately after ROSC
- Outcome: Survival to hospital discharge with good neurologic outcome and survival to hospital discharge were ranked as critical outcomes. ROSC was ranked as an important outcome. For antiarrhythmic drugs after ROSC, rearrest was included as an important outcome.
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) are eligible for inclusion.
- Time frame: All years and languages were included if there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to August 15, 2017.

Treatment Recommendations

We suggest the use of amiodarone or lidocaine in adults with shock-refractory VF/pVT (weak recommendation, low certainty evidence).

We suggest against the routine use of magnesium in adults with shock-refractory VF/pVT (weak recommendation, very low-certainty evidence).

The confidence in effect estimates is currently too low to support an ALS Task Force recommendation about the use of bretylium, nifekalant, or sotalol in the treatment of adults in cardiac arrest with shock refractory VF/pVT.

The confidence in effect estimates is currently too low to support an ALS Task Force recommendation about the use of prophylactic antiarrhythmic drugs immediately after ROSC in adults with VF/pVT cardiac arrest.^{4,5}

IV Versus IO Drug Delivery (ALS 2046: SysRev)

Rationale for Review

This is a new ALS question that was based on the increasing use of IO access during adult resuscitation. It can often be difficult to obtain IV access, especially in the prehospital setting. IO access as an alternative to IV access is increasingly used during cardiac arrest. However, whether drugs are as effective when administered intraosseously versus intravenously is unknown. This 2020 CoSTR is informed by a 2020 SysRev.⁸⁸

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults with cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: Placement of an IO cannula and drug administration through this IO during cardiac arrest
- Comparator: Placement of an IV cannula and drug administration through this IV during cardiac arrest
- Outcome: ROSC, or survival/survival with a favorable neurological outcome at hospital discharge, 30 days, or longer
- Study design: Randomized trials, non-RCTs, and observational studies (cohort studies and case-control studies) comparing IO with IV administration of drugs were included. Randomized trials assessing the effect of specific drugs (ie, epinephrine and amiodarone/lidocaine) in subgroups related to IO versus IV administration were also included. Ecological studies, case series, case reports, reviews, abstracts, editorials, comments, letters to the editor, and unpublished studies were not included. Studies assessing cost-effectiveness were included for a descriptive summary.
- Time frame: The literature search was performed on September 12, 2019, and updated on December 17, 2019, with no date restrictions.
- PROSPERO Registration: CRD42020150877

Consensus on Science

For the important outcome of ROSC, we identified very low-certainty evidence (downgraded for risk of bias and inconsistency) from 4 observational studies^{89–92} including 70 419 adults with OHCA, demonstrating an association of worse outcomes with the use of IO access when compared with IV access (adjusted OR, 0.72 [95% CI, 0.68–0.76]; $P < 0.00001$; absolute risk difference, -6.1% [95% CI, -7.1 to -5.2] or 61 fewer per 1000 cardiac arrests had ROSC with IO access compared with IV access [95% CI, 71 fewer to 52 fewer]).

For the critical outcome of survival to hospital discharge, we identified very low-certainty evidence (downgraded for risk of bias and inconsistency) from 4 observational studies^{89–92} including 70 419 adult OHCA, demonstrating an association of worse outcomes with the use of IO access when compared with IV access (adjusted OR, 0.71 [95% CI, 0.63–0.79]; $P < 0.00001$; absolute risk difference, -2.0% [95% CI, -2.5 to -1.4] or 20 fewer per 1000 cardiac arrests with survival to hospital discharge with use of IO access compared with IV access [95% CI, 25 fewer to 14 fewer]).

For the critical outcome of survival to hospital discharge with a favorable neurological outcome, we identified very low-certainty evidence (downgraded for risk of bias and inconsistency) from 3 observational studies^{89,91,92} including 68 619 adult OHCA, demonstrating an association of worse outcomes with the use of IO access when compared with IV access (adjusted OR, 0.60 [95% CI, 0.52–0.69]; $P < 0.00001$; absolute risk difference, -1.9% [95% CI, -2.3 to -1.5] or 19 fewer per 1000 cardiac arrests with survival to hospital discharge with a favorable neurological outcome with use of IO access compared with IV access [95% CI, 23 fewer to 15 fewer]).

In addition to these findings from observational studies, we identified 2 RCTs of drug administration during cardiac arrest that performed subgroup analyses according to IO versus IV route of administration.^{84,85} None of the comparisons showed statistically significant effect modification. The point estimates generally favored IV access as compared with IO access, except for the outcome of ROSC in the PARAMEDIC2 trial where the effect of epinephrine was similar when given IV or IO. These 2 trials were underpowered to assess such interactions for any outcomes other than ROSC.

Treatment Recommendations

We suggest IV access as compared with IO access as the first attempt for drug administration during adult cardiac arrest (weak recommendation, very low-certainty evidence).

If attempts at IV access are unsuccessful or IV access is not feasible, we suggest IO access as a route for drug administration during adult cardiac arrest (weak recommendation, very low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-2](#). Although the overall certainty in the evidence is very low, the current evidence suggests that outcomes might be better with IV versus IO drug administration. The task force discussed the possibility of unaccounted-for confounders in comparing patients for whom an IV could be obtained with those who required IO placement for access. The task force also discussed that 2015 council guidelines suggest that IO access should be used only if IV access is “difficult or impossible”¹⁷ or “not readily available.”⁹³ The included studies did not enable meaningful analyses of specific subgroups. The documented IO site was primarily tibial, but the site was often not documented. As such, no statements can be made about difference between tibial and humeral (or other) IO sites. All studies were conducted in OHCA patients. Although IHCA patients are likely to have existing IV access, this is not universally true. Although there might be differences in provider skills and patient characteristics between OHCA and IHCA, we consider it unlikely that these would lead to substantial effect modification. As such, the above recommendations apply to both IHCA and OHCA.

Knowledge Gap

- The overall certainty in the evidence is very low. As such, there is clinical equipoise for additional trials related to IV versus IO drug administration during cardiac arrest. These could include trials that directly compare IV to different sites of IO access (eg, tibial, humeral).

Steroids During CPR (ALS 433: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults who are in cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: Corticosteroid or mineralocorticoid administration during CPR
- Comparator: Not using steroids
- Outcome: Survival with favorable neurological/functional outcome at discharge, 30 days, 60 days, 180 days, and/or 1 year; survival only at discharge, 30 days, 60 days, 180 days, and/or 1 year; ROSC
- Intra-arrest steroid use was last reviewed in 2015.^{1,7} The EvUp for intra-arrest steroid use is included in [Supplement Appendix C-6](#). The search identified 2 large, population-based observational studies published since the 2015 CoSTR,^{94,95} both of which suggest a possible association between the use of corticosteroids during CPR and improved survival. Three ongoing clinical trials on this topic were

also identified (NCT02790788, NCT03640949, NCT03317197). The task force will prioritize a SysRev when the results of these trials become available.

Treatment Recommendation

This treatment recommendation (below) is unchanged from 2015.^{1,7}

For IHCA, the task force was unable to reach a consensus recommendation for or against the use of steroids during cardiac arrest.

We suggest against the routine use of steroids during CPR for OHCA (weak recommendation, very low-certainty evidence).

Buffering Agents for Cardiac Arrest (ALS 483: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults with cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: The use of buffering agents alone or combination with other drugs
- Comparator: Not using drugs (or a standard drug regimen)
- Outcome: ROSC, survival, survival with favorable neurological outcome
- This topic was last reviewed in 2010.^{6,8} An EvUp was completed for 2020 and is included in [Supplement Appendix C-7](#). One small RCT and 4 observational studies were identified.⁹⁶⁻¹⁰⁰ An updated SysRev was not considered necessary.

Treatment Recommendation

This treatment recommendation (below) is unchanged from 2010.^{6,8}

Routine administration of sodium bicarbonate for treatment of IHCA and OHCA is not recommended.

Drugs for Torsades de Pointes (ALS 457: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults with torsades de pointes in any setting (in-hospital or out-of-hospital)
- Intervention: Any drug or combination of drugs
- Comparator: Not using drugs or alternative drugs
- Outcome: ROSC, survival, or survival with favorable neurological outcome
- This PICO was last reviewed in 2010.^{6,8} An EvUp is included in [Supplement Appendix C-8](#). No studies meeting inclusion criteria were identified, and thus consideration of an updated SysRev was not suggested.

Treatment Recommendation

This treatment recommendation (below) is unchanged from 2010.^{6,8}

Polymorphic wide-complex tachycardia associated with familial long QT may be treated with IV magnesium, pacing, and/or beta blockers; however, isoprenaline should be avoided.

Polymorphic wide-complex tachycardia associated with acquired long QT may be treated with IV magnesium.

Addition of pacing or IV isoprenaline may be considered when polymorphic wide-complex tachycardia is accompanied by bradycardia or appears to be precipitated by pauses in rhythm.

INTRA-ARREST PROGNOSTICATION

Point-of-Care Echocardiography for Prognostication During CPR (ALS 658: SysRev)

Rationale for Review

In 2015, the question of whether the use of cardiac ultrasound during CPR changed outcomes was reviewed.^{1,7} This question has not been reviewed for the 2020 CoSTR for ALS, and the 2015 CoSTR currently remains: We suggest that if cardiac ultrasound can be performed without interfering with standard advanced cardiovascular life support protocols, it may be considered as an additional diagnostic tool to identify potentially reversible causes (weak recommendation, very low-quality evidence).^{1,7}

The current question is different from that mentioned above and was prioritized by the ALS Task Force due to the increasing popularity of the use of point-of-care echocardiography during cardiac arrest as a prognostic tool, as well as concern about potential pitfalls for misinterpretation of ultrasound findings. A task force–led SysRev of the intra-arrest prognostic capabilities of point-of-care echocardiography was performed to inform the 2020 CoSTR for ALS.¹⁰¹

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults with cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: A particular finding on point-of-care echocardiography during CPR
- Comparator: The absence of that finding or a different finding on point-of-care echocardiography during CPR
- Outcome: Clinical outcomes include, but are not necessarily limited to, ROSC and survival to hospital admission (both important) and the critical outcomes of survival/survival with a favorable neurological outcome at hospital discharge and

survival/survival with a favorable neurological outcome beyond hospital discharge.

- Study design: Randomized trials, non-RCTs, observational studies (cohort studies and case-control studies), registries, and prognosis studies. Ecological studies, case series, case reports, reviews, abstracts, editorials, comments, letters to the editor, or unpublished studies will not be included.
- Time frame: All years and languages were included if there was an English abstract, and there were no date restrictions. The literature search was updated to September 18, 2019.
- PROSPERO Registration: CRD42020150677.

Consensus on Science

The SysRev identified no RCTs and 15 relevant observational studies.^{102–116} The overall certainty of evidence was rated as very low for all outcomes primarily due to risk of bias, inconsistency, and/or imprecision. There was a substantial risk of bias due to prognostic factor measurement, outcome measurement, adjustment for prognostic factors, or confounding. Because of this and a high degree of clinical heterogeneity, no meta-analyses could be performed, and individual studies are difficult to interpret. The consensus on science is summarized in Table 1. The summary of each outcome is separated by the ultrasonographic finding (organized contractility versus nonorganized and/or unspecified motion) and timing of image acquisition (initial, every, any, or subsequent evaluation; or unspecified) because these also varied considerably across studies.

Treatment Recommendation

We suggest against the use of point-of-care echocardiography for prognostication during CPR (weak recommendation, very low-certainty evidence)

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-3](#). This CoSTR specifically addresses the role of ultrasound in prognostication, and in particular prognostication of a favorable outcome that is based on the presence of cardiac motion. In 2015, the task force stated that ultrasound had a potential role in diagnosing reversible causes of cardiac arrest if it could be done without interfering with high-quality CPR, and this recommendation was not reassessed for 2020.^{1,7}

Given the increasing popularity of the use of point-of-care echocardiography for prognostication during attempted resuscitation after cardiac arrest, this comprehensive and rigorous summary of its intra-arrest prognostic capabilities provides valuable information to both the resuscitation science community and bedside clinicians. In making these recommendations, the ALS Task Force considered the following:

Table 1. Estimated Prognostic Test Performance and Prognostic Association for Sonographic Findings on Point-of-Care Echocardiography During Cardiac Arrest to Predict Clinical Outcomes

Outcome	Author, Year	Total Subjects (n), IHCA or OHCA	Sensitivity (Range or 95% CI)	Specificity (Range or 95% CI)	Odds Ratio (Range or 95% CI)
Organized Cardiac Motion (Unspecified Timing of Echocardiography)					
Survival 180 d	Flato, 2015 ¹¹³	49, IHCA	1.0 (95% CI, 0.4–1.0)	0.49 (95% CI, 0.34–0.64)	8.62 (95% CI, 0.44–169.38)
Survival to hospital discharge	Atkinson, 2019 ¹⁰⁸ Flato, 2015 ¹¹³	229, IHCA and OHCA	0.67–1.00	0.51–0.89	13.60–16.63
Survival to hospital admission	Atkinson, 2019 ¹⁰⁸ Blaivas, 2001 ¹⁰⁹	349, OHCA	0.39–1.00	0.91–0.91	6.73–414.56
ROSC	Atkinson, 2019 ¹⁰⁸ Flato, 2015 ¹¹³	229, IHCA and OHCA	0.34–0.79	0.68–0.96	8.07–13.21
Nonorganized and/or Unspecified Cardiac Motion on Initial Echocardiogram					
Good neurological outcome at discharge	Aichinger, 2012 ¹⁰⁷	42, OHCA	1.00 (95% CI, 0.03–1.00)	0.78 (95% CI, 0.62–0.89)	10.26 (95% CI, 0.39–273.09)
Survival to hospital discharge	Gaspari, 2016 ¹¹⁴ Varriale, 1997 ¹⁰⁶ Zengin, 2016 ¹¹⁶	1171,† IHCA and OHCA	0.06–0.91	0.49–0.94	0.38–17.00
Survival to hospital admission	Aichinger, 2012 ¹⁰⁷ Gaspari, 2016 ¹¹⁴ Salen, 2001 ¹⁰⁴ Zengin, 2016 ¹¹⁶	1295,† IHCA and OHCA	0.11–0.92	0.55–0.85	0.75–27.56
ROSC	Gaspari, 2016 ¹¹⁴ Kim, 2016 ¹¹⁵ Varriale, 1997 ¹⁰⁶	861, IHCA and OHCA	0.25–0.64	0.78–1.00	6.33–16.11
Nonorganized and/or Unspecified Cardiac Motion on Every Echocardiogram					
Survival to hospital admission	Aichinger, 2012 ¹⁰⁷ Salen, 2001 ¹⁰⁴	134,* OHCA	0.50–0.80	0.92–1.00	45.33–148.20
Nonorganized and/or Unspecified Cardiac Motion (Unspecified Timing of Echocardiography)					
Good neurological outcome at 180 days	Flato, 2015 ¹¹³	49, IHCA	1.00 (95% CI, 0.40–1.00)	0.49 (95% CI, 0.34–0.64)	8.62 (95% CI, 0.44–169.38)
Good neurological outcome at discharge	Salen, 2005 ¹⁰³	70, OHCA	1.00 (95% CI, 0.03–1.00)	0.86 (95% CI, 0.75–0.93)	17.00 (95% CI, 0.65–446.02)
Survival to hospital discharge	Lien, 2018 ¹⁰²	177, OHCA	0.48 (95% CI, 0.28–0.69)	0.77 (95% CI, 0.69–0.83)	3.09 (95% CI, 1.29–7.37)
Survival to hospital admission	Breitkreutz, 2010 ¹¹⁰ Chua, 2017 ¹¹² Salen, 2001 ¹⁰⁴	291,* OHCA	0.72–0.86	0.60–0.84	9.14–14.00
ROSC	Chardoli, 2012 ¹¹¹ Lien, 2018 ¹⁰² Salen, 2005 ¹⁰³ Tayal, 2003 ¹⁰⁵	317, OHCA	0.62–1.00	0.33–0.98	23.18–289.00
Return of Organized Cardiac Motion on Subsequent Echocardiogram					
Survival to hospital discharge	Varriale, 1997 ¹⁰⁶	20, IHCA	0.50 (95% CI, 0.01–0.99)	0.79 (95% CI, 0.54–0.94)	3.75 (95% CI, 0.19–74.06)
ROSC	Varriale, 1997 ¹⁰⁶	20, IHCA	0.67 (95% CI, 0.22–0.96)	1.00 (95% CI, 0.77–1.00)	52.50 (95% CI, 2.10–1300.33)
Coalescent Echo Contrast (ie, Visible Clotted Intra-Cardiac Blood) After 20–30 min of CPR					
Survival to hospital discharge	Varriale, 1997 ¹⁰⁶	20, IHCA	0.00 (95% CI, 0.00–0.84)	0.45 (95% CI, 0.23–0.68)	0.13 (95% CI, 0.01–3.11)
ROSC	Varriale, 1997 ¹⁰⁶	20, IHCA	0.00 (95% CI, 0.00–0.46)	0.21 (95% CI, 0.05–0.51)	0.02 (95% CI, 0.00–0.53)
Sonographic Evidence of Treatable Pathology					
Survival to hospital discharge	Gaspari, 2016 ¹¹⁴ Varriale, 1997 ¹⁰⁶ Zengin, 2016 ¹¹⁶	1130,† IHCA and OHCA	0.00–0.15	0.89–0.98	1.32–4.25

(Continued)

Table 1. Continued

Outcome	Author, Year	Total Subjects (n), IHCA or OHCA	Sensitivity (Range or 95% CI)	Specificity (Range or 95% CI)	Odds Ratio (Range or 95% CI)
Survival to hospital admission	Zengin, 2016 ¹¹⁶	531,† IHCA and OHCA	0.03–0.04	0.95–0.99	0.61–4.70
ROSC	Chardoli, 2012 ¹¹¹ Lien, 2018 ¹⁰² Tayal, 2003 ¹⁰⁵ Varriale, 1997 ¹⁰⁶	317,† IHCA and OHCA	0.00–1.00	0.84–0.94	0.38–125.00

IHCA indicates in-hospital cardiac arrest; OHCA, out-of-hospital cardiac arrest; and ROSC, return of spontaneous circulation.

*Studies did not report these data for all enrolled subjects; n is lower than the total of all subjects enrolled.

†Gaspari et al and Zengin et al report multiple sonographic findings within a given category on the same subjects; n reflects composite variable “subject-assessments.”

There were inconsistent definitions and terminology about the sonographic evidence of cardiac motion. This included wide variation in the classification of anatomy, type of motion, and timing of point-of-care echocardiogram. The task force encourages the establishment of uniform definitions and terminology to describe sonographic findings of cardiac activity during cardiac arrest.

Most of the identified studies suffer from high risk of bias related to prognostic factor measurement, outcome measurement, lack of adjustment for other prognostic factors, and confounding from self-fulfilling prophecy and unspecified timing of point-of-care echocardiography. Because the risk of bias and heterogeneity across studies was high, no meta-analyses were performed. The evidence supporting use of point-of-care echocardiography as a prognostic tool during cardiac arrest is uniformly of very low certainty. Clinicians should interpret sonographic findings during cardiac arrest in light of these limitations. The task force encourages subsequent investigators studying point-of-care echocardiography during cardiac arrest to identify methodology that mitigates these risks of bias.

Only 2 studies^{113,114} reported estimates of inter-rater reliability (Kappa 0.63 and 0.93). More uniform reporting of inter-rater reliability of point-of-care echocardiography interpretation in future investigations is important.

No sonographic finding had sufficient and/or consistent sensitivity for any clinical outcome for its absence to be used as a sole criterion to stop resuscitation, but the certainty of this evidence is very low.

Some sonographic findings had higher ranges of specificity for clinical outcomes, but the certainty of this evidence is very low.

The impact of ECPR on the prognostic accuracy of point-of-care echocardiography is uncertain.

Point-of-care echocardiography may still be useful to diagnose treatable etiologies of cardiac arrest or to intermittently assess response to resuscitative treatments. These applications are not within the scope of this particular PICOST question. We do, however, caution against overinterpreting the finding of right-ventricular dilation in isolation as a diagnostic indicator of massive pulmonary embolism. Right-ventricular dilation

begins a few minutes after onset of cardiac arrest as blood shifts from the systemic circulation to the right heart along a pressure gradient.^{117,118} Right-ventricular dilation was uniformly observed in a porcine model of cardiac arrest across etiologies of hypovolemia, hyperkalemia, and primary arrhythmia.¹¹⁹

Clinicians should be cautious about potentially prolonging interruptions in chest compressions when using point-of-care echocardiography during cardiac arrest.^{120,121} Several strategies to minimize these interruptions have been proposed.^{122,123}

Point-of-care echocardiography is subject to availability of equipment and skilled operators.

Knowledge Gaps

- There is no standardized or uniform definition of cardiac motion visualized on point-of-care echocardiography during cardiac arrest.
- There are very few prognostic factor studies of point-of-care echocardiography during cardiac arrest performed with methodology that minimizes risk of bias.
- The inter-rater reliability of point-of-care echocardiography during cardiac arrest is uncertain.
- The relative roles and feasibility of transesophageal versus transthoracic echocardiography during CPR require research.

ETCO₂ to Predict Outcome of Cardiac Arrest (ALS 459: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults who are in cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: Any ETCO₂ level value, when present
- Comparator: Any ETCO₂ level below that value
- Outcome: ROSC, survival, survival with favorable neurological outcome
- This topic was last updated in a published 2015 CoSTR,^{1,7} and the SysRev that informed this CoSTR was published in 2018.¹²⁴ The 2 EvUps are included in [Supplement Appendix C-9a and C-9b](#). A search

from December 2013 to November 2019 identified 7 new observational studies^{74,80,125–129} in addition to the previous SysRev.¹²⁴ The task force discussed the low likelihood of an updated SysRev leading to a change in treatment recommendations based on the available studies, and therefore did not prioritize this topic for a SysRev at this time. Future studies and SysRevs should consider trends and changes in ETCO₂ values during CPR in addition to the significance of single ETCO₂ values. The 2015 treatment recommendations remain unchanged.^{1,7}

Treatment Recommendations

This treatment recommendation (below) is unchanged from 2015.^{1,7}

We recommend against using ETCO₂ cutoff values alone as a mortality predictor or for the decision to stop a resuscitation attempt (strong recommendation, low-quality evidence).

We suggest that an ETCO₂ of 10 mmHg or greater measured after tracheal intubation or after 20 minutes of resuscitation may be a predictor of ROSC (weak recommendation, low-quality evidence).

We suggest that an ETCO₂ of 10 mmHg or greater measured after tracheal intubation, or an ETCO₂ of 20 mmHg or greater measured after 20 minutes of resuscitation, may be a predictor of survival to discharge (weak recommendation, moderate-quality evidence).

CARDIAC ARREST IN SPECIAL CIRCUMSTANCES

Cardiac Arrest Associated With Pulmonary Embolism (ALS 435, 581: SysRev)

Rationale for Review

Pulmonary embolism (PE) is a potentially reversible cause of cardiac arrest. Whether chances for ROSC and survival may be significantly higher if a PE is present and can be treated is not well established because research has been limited to-date. This topic was last reviewed in 2015.^{1,7} The specific role of ECPR was not addressed in this updated SysRev because ECPR was addressed in the previous 2019 CoSTR summary.^{2,3} The role of ECPR for the treatment of PE and cardiac arrest is discussed in the justification section that follows.

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults in cardiac arrest due to PE or suspected PE in any setting (in-hospital or out-of-hospital)
- Intervention: Any specific alteration in the ALS treatment algorithm (eg, fibrinolytics or any other)
- Comparator: Standard ALS care

- Outcome: Survival with favorable neurological outcome at discharge, 30 days, 60 days, 180 days, and/or 1 year; survival at discharge, 30 days, 60 days, 180 days, and/or 1 year (all critical); ROSC (important)
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) are eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) are excluded.
- Time frame: All years and languages were included if there was an English abstract. Literature search was updated to October 2019.
- PROSPERO Registration: Registered with ILCOR Science Advisory Committee October 6, 2019. This SysRev was done as an update of the 2015 CoSTR SysRev and PROSPERO registration was not done.

Consensus on Science

Fibrinolysis

For the critical outcome of survival with favorable neurological outcome at 30 days, we identified very low-certainty evidence (downgraded for serious imprecision and very serious risk of bias) from a subgroup of 37 patients with confirmed PE from 1 RCT comparing fibrinolytics with placebo during cardiac arrest¹³⁰ finding no difference between groups [tenecteplase 2/15, (13.3%) versus placebo, 0/22 (0%)] [risk ratio (RR), 7.19; 95% CI, 0.37–139.9]. We also identified very low-certainty evidence (downgraded for risk of bias) from a single observational study that found no difference (10% with fibrinolysis versus 5% without; adjusted RR [ARR], 1.97; 95% CI, 0.70–5.56).¹³¹

For the critical outcome of survival at 30 days, very low-certainty evidence (downgraded for risk of bias) from one observational study showed an association between improved survival and administration of fibrinolysis (16% with fibrinolysis versus 6% without; $P=0.005$).¹³¹

For the critical outcome of survival to hospital discharge, very low-certainty evidence (downgraded for very serious risk of bias and imprecision) from 3 retrospective observational studies showed no association between administration of fibrinolysis and survival (10.5% survival with fibrinolysis versus 8.7% without;¹³² 9.5% survival with fibrinolysis versus 4.8% without¹³³ and 19.4% survival with fibrinolysis versus 6.7% without (RR, 2.9; 95% CI, 0.75–13.8).¹³⁴

For the important outcome of ROSC, very low-certainty evidence from 1 study (downgraded for very serious risk of bias) showed benefit associated with the use of fibrinolytic drugs compared with no fibrinolytic drugs in patients with PE (81.0% with fibrinolysis versus 42.9% without; $P=0.03$).¹³³ Two other studies provided very low-certainty evidence (downgraded for very serious risk of bias) of no difference in ROSC (66.7% with fibrinolysis versus 43.3% without [RR, 1.5; 95%

CI, 0.8–8.6] and 47.4% with fibrinolysis versus 47.8% without; $P=0.98$).^{132,134}

For the outcome of survival at 24 hours, very low-certainty evidence (downgraded for risk of bias) from 1 observational study showed no difference associated with fibrinolysis (66% with fibrinolysis versus 63% without; $P=0.76$),¹³¹ whereas another study showed benefit associated with fibrinolysis (52.8% with fibrinolysis versus 23.3% without; RR, 2.3; 95% CI, 1.1–4.7).¹³⁴

Surgical Embolectomy

We found very low-certainty evidence (downgraded for very serious risk of bias) from 2 case series^{135,136} with no control groups and a total of 21 patients requiring CPR with a 30-day survival rate of 12.5% and 71.4%, respectively.

Percutaneous Mechanical Thrombectomy

For the important outcome of ROSC, very low-certainty evidence (downgraded for very serious risk of bias and very serious imprecision) from 1 case series of 7 patients with cardiac arrest with no control group,¹³⁷ ROSC was achieved in 6 of 7 patients (85.7%) treated with percutaneous mechanical thrombectomy.

The overall certainty of evidence was rated as very low primarily due to a very serious risk of bias and inconsistency. For this reason, no meta-analyses were performed.

Treatment Recommendations

These recommendations (below) are unchanged from 2015.^{1,7} We suggest administering fibrinolytic drugs for cardiac arrest when PE is the suspected cause of cardiac arrest (weak recommendation, very low-certainty evidence).

We suggest the use of fibrinolytic drugs or surgical embolectomy or percutaneous mechanical thrombectomy for cardiac arrest when PE is the known cause of cardiac arrest (weak recommendation, very low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-4](#). The task force considered that mechanical or surgical thrombectomy would be used only if the patient had a confirmed PE. No RCTs were identified and no meta-analysis was undertaken given the limited evidence.

The task force considered that 2% to 7% of patients with OHCA have a PE,^{130,131} and this figure is probably higher for patients with IHCA. The task force acknowledged that ECPR techniques are now frequently used in patients with cardiac arrest from suspected PE in those settings where it is feasible. This role of ECPR for advanced life support was addressed by the 2019 CoSTR summary, and the considered studies included patients with PE.^{2,3} Specifically in patients with PE, ECPR may potentially

facilitate the use of fibrinolysis or mechanical or surgical embolectomy, but the evidence is of very low certainty.

The task force considered the increased risk of bleeding from fibrinolysis if it is administered to patients without PE. The Thrombolysis in Cardiac Arrest (TROICA) Study—which is the largest study of thrombolysis during cardiac arrest—showed an increased risk of bleeding in the thrombolysis group (any intracranial hemorrhage, 2.7% versus 0.4%; RR, 6.95 [95% CI, 1.59–30.41]; $P=0.006$), but major bleeding complications did not occur more often (symptomatic intracranial hemorrhage, 0.8% versus 0%; RR, 8.93 [95% CI, 0.48–165.45]; $P=0.13$; major nonintracranial hemorrhage, 7.7% versus 6.4%; RR, 1.21 [95% CI, 0.77–1.88]; $P=0.48$; ischemic stroke, 0.8% versus 0.6%; RR, 1.32 [95% CI, 0.30–5.88]; $P=1.00$).¹³⁰ Patients are far more likely to die from the cardiac arrest than from the treatment.

Knowledge Gap

- Optimal drug and dosing strategy for fibrinolysis during CPR with a suspected or actual PE
- Intra-arrest prediction of PE during CPR

Cardiac Arrest in Pregnancy (ALS 436: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Pregnant women who are in cardiac arrest in any setting
- Intervention: Any specific intervention(s)
- Comparator: Standard care (usual resuscitation practice)
- Outcome: ROSC; survival to discharge, 30 days, or longer; survival with favorable neurological outcome at discharge, 30 days, or longer
- An EvUp is included in [Supplement Appendix C-10](#). Since the 2015 CoSTR,^{1,7} 7 new small observational studies were identified, 5 of which focused on association of timing of delivery with outcome of cardiac arrest and other factors associated with maternal and fetal mortality.^{138–142} Due to the very small size of most studies, an updated SysRev was not suggested. The 2015 treatment recommendation remains unchanged.^{1,7}

Treatment Recommendation

This treatment recommendation (below) is unchanged from 2015.^{1,7}

We suggest delivery of the fetus by perimortem cesarean delivery for women in cardiac arrest in the second half of pregnancy (weak recommendation, very low-quality evidence).

There is insufficient evidence to define a specific time interval by which delivery should begin.

High-quality usual resuscitation care and therapeutic interventions that target the most likely cause(s) of cardiac arrest remain important in this population.

There is insufficient evidence to make a recommendation about the use of left-lateral tilt and/or uterine displacement during CPR in the pregnant patient.

Opioid Toxicity (ALS 441: EvUp)

Death from opioid toxicity is an important public health issue in many countries. The issue of first aid education for opioid toxicity has been addressed by the EIT Task Force ScopRev 891.^{142a,142b}

Population, Intervention, Comparator, and Outcome

- Population: Adults who are in cardiac arrest or respiratory arrest due to opioid toxicity in any setting (in-hospital or out-of-hospital)
- Intervention: Any specific therapy (eg, naloxone, bicarbonate, or other drugs)
- Comparator: Usual ALS care
- Outcome: Survival with favorable neurological/functional outcome at discharge, 30 days, 60 days, 180 days, and/or 1 year; survival only at discharge, 30 days, 60 days, 180 days, and/or 1 year; ROSC were defined as critical or important outcomes
- This topic was last reviewed in 2015.^{1,7} The EvUp is included in [Supplement Appendix C-11](#). The search was conducted for studies published from September 1, 2013, to September 13, 2019. There was insufficient evidence to support consideration of a stand-alone ALS SysRev, but an updated SysRev with other task forces was suggested.

Treatment Recommendations

This treatment recommendation (below) is unchanged from 2015.^{1,7}

We recommend the use of naloxone by IV, intramuscular, subcutaneous, IO, or intranasal routes in respiratory arrest associated with opioid toxicity (strong recommendation, very low-quality evidence). The dose of naloxone required will depend on the route.

We can make no recommendation about the modification of standard ALS in opioid-induced cardiac arrest.

POSTRESUSCITATION CARE

The last update of postresuscitation care was published in the 2015 CoSTR.^{1,7} Since that publication, there have been many reported studies of postresuscitation care.¹⁴³

Oxygen Dose After ROSC in Adults (ALS 448: SysRev)

Rationale for Review

Both hypoxemia and hyperoxemia during postresuscitation care have been associated with worse outcomes. Hypoxemia may worsen ischemic brain injury and injury to other organs, and hyperoxemia may lead to increased

oxidative stress and organ damage after reperfusion. A SysRev was conducted to inform this 2020 CoSTR for ALS.¹⁴⁴

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Unresponsive adults with sustained ROSC after cardiac arrest in any setting
- Intervention: A ventilation strategy targeting a specific oxygen saturation and/or Pao₂
- Comparator: Treatment without specific targets or with an alternate target to the intervention
- Outcome: Critical outcomes include survival/survival with a favorable neurological outcome at hospital discharge or 30 days; and survival/survival with a favorable neurological outcome after hospital discharge or 30 days (eg, 90 days, 180 days, 1 year).
- Study design: Randomized trials, non-RCTs, and observational studies (cohort studies and case-control studies) with a control group (ie, patients treated with no specific oxygen saturation and/or Pao₂ targets or an alternative target to the intervention) were included. Animal studies, ecological studies, case series, case reports, reviews, abstracts, editorials, comments, and letters to the editor were not included. There were no limitations on publication period or study language, if there was an English abstract. The population included patients with IHCA or OHCA of any origin. Unpublished studies (eg, conference abstracts, trial protocols) were excluded. The cited SysRev¹⁴⁴ was performed without age restriction, and the evidence from adult studies (generally defined as older than 16 years or 18 years or older) is included here.
- Time frame: All years and languages were included. The literature search was updated to August 22, 2019.
- PROSPERO Registration: CRD42020150877

Consensus on Science

The evidence from the 6 RCTs identified in the SysRev is summarized in Table 2. Trials generally failed to show a benefit of a titrated (lower percentage of oxygen) approach compared with standard care (higher percentage of oxygen). A subgroup analysis of postresuscitation patients in one larger RCT, however, found better survival in patients for whom hyperoxemia was aggressively avoided.¹⁴⁵ In addition, results from 10 observational studies rated as having only serious risk of bias were inconsistent. Four^{146–149} found an association between hyperoxemia (variable definitions, but most often Pao₂ greater than 300 mmHg) and either worse survival or worse survival with neurological outcome, whereas the other 6^{150–155} found

Table 2. Overview of Included Randomized Trials

Study, Year	Country	Years of Inclusion	Number of Patients	Intervention	Comparator	Relative Risk [95% CI]	Absolute Risk Reduction [95% CI]	Certainty
Favorable Neurological Outcome at 6 mo*—ICU Initiation								
Jakkula, 2018 ¹⁵⁶	Finland and Denmark	2016–2017	120	Normoxia for 36 h (10–15 kPa)	Moderate hyperoxia for 36 h (20–25 kPa)	1.13 [0.87–1.47]	79 more per 1000 [79 fewer to 287 more]	Moderate†
Mackle, 2019 ¹⁴⁵	Australia and New Zealand	2015–2018	164	Conservative oxygen (O ₂ Sat 90%–97%)	Standard oxygen (O ₂ Sat >90%)	1.40 [0.93–2.13]	128 more per 1000 [22 fewer to 361 more]	Very low‡
Survival to Hospital Discharge With Favorable Neurological Outcome#—Prehospital Initiation								
Young, 2014 ¹⁵⁷	New Zealand	2012–2013	17	Titration oxygen for 72 h (O ₂ Sat 90%–94%)	Standard oxygen for 72 h (O ₂ Sat >95%)	0.56 [0.14–2.29]	196 fewer per 1000 [382 fewer to 573 more]	Very low¶
Kuisma, 2006 ¹⁵⁸	Finland	Not recorded	28	Low oxygen prehospital (30%)	High oxygen prehospital (100%)	1.33 [0.63–2.84]	141 more per 1000 [159 fewer to 789 more]	Low§
Survival to Hospital Discharge—ICU Initiation								
Jakkula, 2018 ¹⁵⁶	Finland and Denmark	2016–2017	120	Normoxia for 36 h (10–15 kPa)	Moderate hyperoxia for 36 h (20–25 kPa)	1.07 [0.84–1.36]	46 more per 1000 [106 fewer to 238 more]	Moderate§
Survival to 90 Days—ICU Initiation								
Mackle, 2019 ¹⁴⁵	Australia and New Zealand	2015–2018	164	Conservative oxygen (O ₂ Sat 90%–97%)	Standard oxygen (O ₂ Sat >90%)	1.39 [1.01–1.92]	160 more per 1000 [4 more to 377 more]	Low††
Survival to Hospital Discharge—Prehospital Initiation								
Meta-analysis Kuisma 2006 ¹⁵⁸ /Bray, 2018 ¹⁵⁹	Finland, Australia	Not recorded, ¹⁵⁸ 2015–2017 ¹⁵⁹	89	Low oxygen prehospital (30% or 2–4 L/min)	High oxygen prehospital (100% or ≥10 L/min)	0.97 [0.68–1.37]	18 fewer per 1000 [194 fewer to 224 more]	Very low§
Thomas, 2019 ¹⁶⁰	United Kingdom	2014–2015	35	Titration oxygen prehospital (O ₂ Sat 94%–98%)	Standard oxygen prehospital (O ₂ Sat 100%)	3.15 [1.04–9.52]	379 more per 1000 [7 more to 1000 more]	Very low¶
Young,** 2014 ¹⁵⁷	New Zealand	2012–2013	17	Titration oxygen for 72 h (O ₂ Sat 90%–94%)	Standard oxygen for 72 h (O ₂ Sat >95%)	1.13 [0.41–3.08]	58 more per 1000 [262 fewer to 924 more]	Very low¶

ICU indicates intensive care unit; and Sat, saturation.

*Defined as either Cerebral Performance Category (CPC) 1–2 or Extended Glasgow Outcome Score of 5–8.

†Downgraded for imprecision.

‡Downgraded for risk of bias and imprecision.

§Downgraded 2 levels for imprecision.

¶Downgraded for indirectness and 2 levels for imprecision.

#Defined as CPC 1–2 or discharge to home.

**Intervention initiated prehospital but continued after admission.

††Downgraded 2 levels for risk of bias.

no such association. Hypoxemia was found to be associated with worse outcome in adjusted analysis in 1 of these studies.¹⁴⁹

Treatment Recommendations

We suggest the use of 100% inspired oxygen until the arterial oxygen saturation or the partial pressure of arterial oxygen can be measured reliably in adults with ROSC after cardiac arrest in any setting (weak recommendation, very low-certainty evidence).

We recommend avoiding hypoxemia in adults with ROSC after cardiac arrest in any setting (strong recommendation, very low-certainty evidence).

We suggest avoiding hyperoxemia in adults with ROSC after cardiac arrest in any setting (weak recommendation, low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-5](#). In making the recommendation to avoid hypoxemia, the task force acknowledges that the evidence is of very low certainty. The task force concluded that the known harm that can result from hypoxia justifies its avoidance, and detection of hypoxemia may be the best surrogate for or precursor of tissue hypoxia. The

Downloaded from <http://ahajournals.org> by on October 21, 2020

suggestion to avoid hyperoxemia is based on low- to moderate-certainty evidence that showed either harm or no benefit from hyperoxemia. The definitions used for hyperoxemia varied, ranging from an oxygen saturation greater than 97% measured by pulse oximeter to a P_{aO_2} up to 20 to 25 kPa (150–188 mmHg) in the available RCTs. In light of the possible benefit and lack of evidence for harm, the task force suggests targeting normoxemia and avoiding hyperoxemia. The task force acknowledges that the primary randomized trial evidence suggesting benefit from avoiding hyperoxemia is from a subgroup analysis only, and data from the 3 ongoing trials (NCT03138005, NCT03653325, NCT03141099) will be helpful.

The task force felt that titration of oxygen should not be attempted until oxygen levels (peripheral oxygen saturation or partial pressure of oxygen in arterial blood) could be measured reliably. Some of the randomized trials conducted in the prehospital setting, although very small, reported more desaturation of arterial blood in the lower oxygen group, which reinforces the task force suggestion to administer 100% oxygen until reliable measurement of oxygen level is possible. This is likely to be more important in the prehospital setting.

Knowledge Gap

- Randomized trials comparing lower oxygen target strategies with higher oxygen target strategies or usual care in postarrest patients have thus far been small and therefore inconclusive. More trials are needed, and 3 trials are underway currently (NCT03138005, NCT03653325, NCT03141099).

Ventilation Strategy After ROSC in Adults (ALS 571: SysRev)

Rationale for Review

Hypocapnia causes cerebral vasoconstriction and hypercapnia leads to cerebral vasodilation. Exactly how variations in P_{aCO_2} affect intracranial pressure and perfusion in the brains of postarrest patients, and whether this affects outcome, remains unclear.¹⁶¹ This topic was last reviewed in 2015.^{1,7} A SysRev¹⁴⁴ was conducted to inform this 2020 ALS CoSTR.

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Unresponsive adults with sustained ROSC after cardiac arrest in any setting
- Intervention: A ventilation strategy targeting a specific P_{aCO_2}
- Comparator: Treatment without specific targets or with an alternate target to the intervention
- Outcome: Critical outcomes include survival/survival with a favorable neurological outcome at hospital discharge or 30 days; and survival/survival with a favorable neurological outcome after

hospital discharge or 30 days (eg, 90 days, 180 days, 1 year).

- Study design: Randomized trials, non-RCTs, and observational studies (cohort studies and case-control studies) with a control group (ie, patients treated with no specific P_{aCO_2} targets or an alternative target to the intervention) were included. Animal studies, ecological studies, case series, case reports, reviews, abstracts, editorials, comments, and letters to the editor were not included. There were no limitations on publication period or study language, if there was an English abstract. The population included patients with IHCA or OHCA of any origin. Unpublished studies (eg, conference abstracts, trial protocols) were excluded. The cited SysRev¹⁴⁴ was done without age restriction, and the evidence from adult studies (generally defined as older than 16 years or 18 years or older) is included here.
- Time frame: All years and languages were included. The literature search was updated to August 22, 2019.
- PROSPERO Registration: CRD42020150877

Consensus on Science

The task force concluded that differences in the P_{aCO_2} targets used in the arms of the 2 RCTs identified^{156,162} precluded meta-analysis.

For the critical outcome of favorable neurological outcome (defined as CPC 1–2) at 6 months, we identified low-certainty evidence from 1 RCT enrolling 120 patients and comparing a ventilation strategy targeting high-normal P_{aCO_2} (5.8–6.0 kPa/43.5–45 mmHg) with one targeting low-normal P_{aCO_2} (4.5–4.7 kPa/33.7–35.2 mmHg) and failing to show benefit from the higher P_{aCO_2} strategy (RR, 0.84; 95% CI, 0.64–1.10; ARR, 113 fewer per 1000; 95% CI, from 254 fewer to 70 more).¹⁵⁶ For the critical outcome of favorable neurological outcome (defined as an extended Glasgow Outcomes Scale ≥ 5) at 6 months, we identified low-certainty evidence (downgraded for inconsistency and imprecision) from 1 RCT enrolling 83 patients and comparing a ventilation strategy targeting moderate hypercapnia (P_{aCO_2} 50–55 mmHg/6.7–7.3 kPa) with one targeting normocapnia (P_{aCO_2} 35–45 mmHg/4.7–6.0 kPa) and failing to show benefit from the higher P_{aCO_2} strategy (RR, 1.28; 95% CI, 0.83–1.96; ARR, 129 more per 1000; 95% CI, from 78 fewer to 443 more).¹⁶²

For the critical outcome of survival to 30 days we identified low-certainty evidence (downgraded for inconsistency and imprecision) from 1 RCT enrolling 120 patients and comparing a ventilation strategy targeting high-normal P_{aCO_2} (5.8–6.0 kPa/43.5–45 mmHg) with one targeting low-normal P_{aCO_2} (4.5–4.7 kPa/33.7–35.2 mmHg) and failing to show benefit from the higher

Paco₂ strategy (RR, 0.81; 95% CI, 0.63–1.05; ARR, 143 fewer per 1000; 95% CI, from 279 fewer to 38 more).¹⁵⁶

For the critical outcome of survival to discharge we identified low-certainty evidence (downgraded for inconsistency and imprecision) from 1 RCT enrolling 83 patients and comparing a ventilation strategy targeting moderate hypercapnia (Paco₂, 50–55 mmHg/6.7–7.3 kPa) with one targeting normocapnia (Paco₂, 35–45 mmHg/4.7–6.0 kPa) and failing to show benefit from the higher Paco₂ strategy (RR, 1.16; 95% CI, 0.87–1.56; ARR, 101 more per 1000; 95% CI, from 82 fewer to 355 more).¹⁶²

Results were inconsistent across the 6 observational studies rated as having less than critical risk of bias. Hypercapnia was associated with improved outcomes in 2 studies^{155,163} and worse outcomes in 2 studies.^{149,164} There was no association between hypercapnia and outcomes in the remaining 2 studies.^{152,165} Results were similar for hypocapnia although no studies found an association with improved outcomes.

Treatment Recommendations

There is insufficient evidence to suggest for or against targeting mild hypercapnia compared with normocapnia in adults with ROSC after cardiac arrest.

We suggest against routinely targeting hypocapnia in adults with ROSC after cardiac arrest (weak recommendation, low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-6](#). Evidence from existing randomized trials and observational studies is very inconsistent. Both randomized trials failed to show any effect from different Paco₂ targets, but the trials were small and used different target ranges, precluding meta-analysis. Observational studies were evenly distributed in showing benefit, harm or no effect associated with hypercapnia. Hypocapnia results were also inconsistent, although no studies found an association with benefit. In light of the lack of evidence for benefit, and lack of consistent evidence for harm from Paco₂ levels higher than normal, the task force did not think there was sufficient evidence to suggest for or against targeting mild hypercapnia compared with normocapnia. An ongoing trial investigating this comparison may bring clarity to this issue (NCT03114033).

For hypocapnia, very limited evidence suggests either no benefit or harm, supporting the task force's suggestion against targeting hypocapnia.

Although the task force discussed whether patients with baseline chronic lung disease and chronic CO₂ retention might respond differently to different Paco₂ targets, no evidence addressing this subgroup was found. The task force agreed it would be reasonable to adjust Paco₂ targets in patients with known chronic CO₂

retention, but this is expert opinion only because no evidence was identified on this topic.

The prior treatment recommendation (2015^{1,7}) was a suggestion to maintain normocapnia. The updated treatment recommendation allows for continuing this approach, while emphasizing that we do not currently know if targeting normocapnia is beneficial, harmful, or equal in comparison to targeting hypercapnia. The task force discussed the possible complication of acidemia from hypercapnia. The presence or absence of metabolic acidosis requires consideration when choosing a ventilation strategy and Paco₂ target, and metabolic acidosis is common in postarrest patients. The Paco₂ targets or ranges also differed somewhat across studies. For this reason, the task force chose not to define specific numeric targets because no optimal target or range has been made clear. Additionally, opinions vary on whether arterial blood gas analysis in patients receiving targeted temperature management (TTM) should be adjusted for temperature. Once again, trials differed in their approach. Approaches to blood gas interpretation in regard to temperature also varied across the observational studies. These variations in methodology and in definitions of target ranges prohibit the task force from being able to recommend specific numbers or a specific method for blood gas analysis for systems implementing these recommendations.

Knowledge Gaps

- Randomized trials comparing strategies targeting mild hypercapnia with strategies targeting normocapnia have thus far been small and therefore inconclusive. A much larger randomized trial is currently underway (NCT03114033).
- How Paco₂ targets should be adjusted in those with chronic CO₂ retention is unknown.

Postresuscitation Hemodynamic Support (ALS 570: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adults with ROSC after cardiac arrest in any setting
- Intervention: Titration of therapy to achieve a specific hemodynamic goal (eg, mean arterial pressure greater than 65 mmHg)
- Comparator: No hemodynamic goal
- Outcome: Any clinical outcome
- An EvUp for this topic was performed and is included in [Supplement Appendix C-12](#). Two RCTs completed since 2015^{156,166} did not find that targeting a specific mean arterial pressure affected outcome, although the studies were not powered for clinical outcomes of survival or neurological outcome. In the absence of ongoing RCTs, and

controversy about the targeting of higher blood pressure, the task force suggests that this topic be considered for a SysRev.

Treatment Recommendations

This treatment recommendation (below) is unchanged from 2015.^{1,7}

We suggest hemodynamic goals (eg, mean arterial pressure, systolic blood pressure) be considered during postresuscitation care and as part of any bundle of postresuscitation interventions (weak recommendation, low-certainty evidence).

There is insufficient evidence to recommend specific hemodynamic goals; such goals should be considered on an individual patient basis and are likely to be influenced by post–cardiac arrest status and preexisting comorbidities (weak recommendation, low-certainty evidence).

Postresuscitation Steroids (ALS 446: EvUp)

Population, Intervention, Comparator, and Outcome

- Population: Adult patients with ROSC after cardiac arrest (prehospital or in-hospital)
- Intervention: Treatment with corticosteroids
- Comparator: Standard care
- Outcome: Survival to hospital discharge with good neurological outcome or survival to hospital discharge (\pm time to shock reversal/shock reversal)
- The 2010 CoSTR addressed steroid use both intra-arrest and postresuscitation.^{6,8} The 2015 CoSTR included only intra-arrest steroid use.^{1,7} The EvUp for postresuscitation steroid use is included in [Supplement Appendix C-13](#). Three small RCTs and a large observational study were identified.^{94,167–169} Two of the RCTs used steroids both during CPR and after ROSC.^{167,168} One recently completed trial that is not yet published was also identified (NCT02790788). The task force recommends a SysRev be undertaken once the recently completed trial is published.

Treatment Recommendations

This treatment (below) is unchanged from 2010.^{6,8}

There is insufficient evidence to support or refute the use of corticosteroids for patients with ROSC after cardiac arrest.

Prophylactic Antibiotics After Cardiac Arrest (ALS 2000: SysRev)

Rationale for Review

This is a new topic prioritized by the ALS Task Force. Infective complications are common in patients admitted to intensive care units (ICUs). After cardiac arrest, pneumonia has been reported in 50% to 60% of patients,^{170,171} which is thought to result in part from

aspiration during the cardiac arrest and resuscitation. In these patients, early and accurate identification of infection is challenging. Standard criteria for identifying infection are affected by patient treatment (ie, TTM) and the pathophysiology of the post–cardiac arrest syndrome (ie, including the systemic inflammatory response). The decision to treat a possible infection needs to be balanced by the need for prudent antibiotic administration to avoid antibiotic resistance. This new topic was prioritized by the ALS Task Force due to the recent publication of a SysRev on the topic.¹⁷² The published SysRev was updated by using the ADOLOPMENT process.¹⁷³

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adult patients after ROSC from cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: Early/prophylactic administration of antibiotics
- Comparator: Delayed/clinically driven administration
- Outcome: Survival or survival with good neurological outcome at hospital discharge or longer (critical), and important outcomes of critical care length of stay, infective complications, or duration of mechanical ventilation
- Study design: Observational and interventional studies if they compared the effect of administration of early or prophylactic antibiotics with delayed or clinically driven administration of antibiotics in adult patients after cardiac arrest. All study types that included a control group were included. Case reports and case series were not eligible for inclusion. There was no restriction on language.
- Time frame: There was no restriction on publication date, and the literature search was completed/updated in October 2019.
- PROSPERO Registration: CRD42016039358 for the original SysRev.¹⁷²

Consensus on Science

For the critical outcome of survival with favorable neurological outcome at ICU discharge or 30 days, we identified low-certainty evidence (downgraded for serious risk of bias and serious imprecision) from 2 RCTs^{171,174} enrolling 254 patients, which showed no benefit of early/prophylactic antibiotic administration (RR, 0.89; 95% CI, 0.71–1.12; $P=0.31$; risk difference, -0.06 ; 95% CI, -0.19 to 0.06 ; $P=0.30$).

For the critical outcome of survival at ICU discharge or 30 days, we identified low-certainty evidence (downgraded for serious risk of bias and serious imprecision) from 2 RCTs^{171,174} enrolling 254 patients, which showed no benefit (RR, 0.95; 95% CI, 0.79–1.14; $P=0.60$; risk difference, -0.03 ; 95% CI, -0.15 to 0.08 ; $P=0.58$). We also identified very low-certainty evidence (downgraded for serious indirectness) from 2 observational

studies. One study¹⁷⁵ enrolling 1604 patients showed no benefit associated with early or prophylactic antibiotic administration compared with delayed/clinically driven administration (OR, 1.16; 95% CI, 0.94 to 1.13; $P=0.18$). The second observational study¹⁷⁶ enrolling 138 patients showed a benefit (data presentation precludes reporting of OR, $P=0.01$).

For the important outcome of infective complications (pneumonia) we identified low-certainty evidence (downgraded for serious risk of bias and serious imprecision) from 2 RCTs^{171,174} enrolling 254 patients, which showed no benefit (RR, 0.75; 95% CI, 0.43 to 1.32; $P=0.32$; risk difference, -0.12 ; 95% CI, -0.23 to 0.00 ; $P=0.05$). There were differences between the studies in methods used to diagnose pneumonia. We found very low-certainty evidence (downgraded for serious risk of bias, serious indirectness, and serious imprecision) from 2 observational studies^{175,177} enrolling 2245 patients, which showed no association between early/prophylactic administration compared with delayed/clinically driven administration (OR, 0.61; 95% CI, 0.61 to 1.62; $P=0.98$). These studies, too, differed in methods used to diagnose pneumonia.

For the important outcome of critical care length of stay we identified low-certainty evidence (downgraded for serious risk of bias and serious imprecision) from 2 RCTs^{171,174} enrolling 248 patients, which showed no benefit (mean difference, 0.47 days; 95% CI, -1.31 to 2.24 ; $P=0.61$).

For the important outcome of duration of mechanical ventilation we identified very low-certainty evidence (downgraded for very serious risk of bias and serious imprecision) from 1 RCT¹⁷⁴ enrolling 60 patients, which showed no benefit (mean difference, 0.20 days; 95% CI, -1.53 to 1.93 ; $P=0.82$).

Treatment Recommendation

We suggest against the use of prophylactic antibiotics in patients after ROSC (weak recommendation, low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-7](#). Meta-analyses of both randomized trials and observational studies showed no overall benefit in the use of prophylactic antibiotics during post-cardiac arrest care. The task force did review the findings of 1 RCT at low overall risk of bias that reported reduced incidence of early pneumonia in patients treated with prophylactic antibiotics.¹⁷¹ Although this study demonstrated the potential efficacy of prophylactic antibiotics, there was no improvement in other clinical outcomes, such as survival or critical care length of stay. Pneumonia affects approximately 50% of ICU patients after cardiac arrest, but this is unlikely to contribute to mortality because most deaths are attributable to neurological failure, cardiovascular failure, or multiorgan failure.^{170,171} A strategy

of prophylactic antibiotic use would likely expose a large number of patients to antibiotics with no specific benefit and increase the risk of development of resistant organisms. The decision to administer antibiotics after cardiac arrest, particularly in the context of gastric aspiration, is challenging and clinicians may have different clinical thresholds for prescribing antibiotics. We did not identify any RCTs enrolling patients after IHCA.

Knowledge Gaps

- Studies of post-ROSC antibiotics after IHCA.
- RCTs that evaluate this question in patients treated with TTM at temperatures other than 32°C to 34°C.
- RCTs powered to determine the effect of prophylactic antibiotics on outcomes such as critical care length of stay or duration of mechanical ventilation.

Post-Cardiac Arrest Seizure Prophylaxis and Treatment (ALS 431, 868: SysRev)

Rationale for Review

Hypoxic-ischemic brain injury is a common cause of death in comatose cardiac arrest survivors. Clinical convulsions and epileptiform activity in the electroencephalogram (EEG) are common, with substantial overlap and an approximate incidence of 20% to 30%.^{178–181} The prognosis for patients with clinical and electrographic seizures is usually poor, but some patients recover and may ultimately have a good neurological outcome.^{180,181} This CoSTR is based on an update of the 2015 SysRev and CoSTR^{1,7} for seizure prophylaxis and treatment in cardiac arrest survivors.

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Unresponsive adults (older than 18 years) with sustained ROSC after cardiac arrest in any setting (in-hospital or out-of-hospital)
- Intervention: One strategy for seizure prophylaxis or treatment
- Comparator: Another strategy or no seizure prophylaxis or treatment
- Outcome: Survival with favorable neurological/functional outcome at discharge, 30 days, 60 days, 180 days, and/or 1 year; survival at discharge, 30 days, 60 days, 180 days, and/or 1 year (all critical); and the important outcome of seizure incidence during index hospitalization (for seizure prophylaxis only)
- Study design: RCTs and nonrandomized studies (non-RCTs, interrupted time series, controlled before-and-after studies, cohort studies) are eligible for inclusion. Unpublished studies (eg, conference abstracts, trial protocols) are excluded.

- Time frame: All years and languages were included if there was an English abstract; unpublished studies (eg, conference abstracts, trial protocols) were excluded. The literature search was updated to September 26, 2019.
- PROSPERO Registration: Registered with ILCOR Science Advisory Committee October 3, 2019. This SysRev was done as an update of the 2015 CoSTR SysRev and PROSPERO registration was not done.

Consensus on Science

Post-Cardiac Arrest Seizure Prophylaxis

For the critical outcomes of survival with favorable neurological outcome to discharge/30 days or longer, and survival to discharge/30 days or longer, 2 prospective RCTs involving a total of 562 subjects provided very low-certainty evidence (downgraded for risk of bias, indirectness and imprecision)^{182,183} of no benefit from seizure prophylaxis. One nonrandomized prospective clinical trial with 107 subjects that used historic controls provided very low-certainty evidence (downgraded for risk of bias, indirectness, and imprecision) of no benefit.¹⁸⁴

For the important outcome of seizure prevention, we identified very low-certainty evidence (downgraded for risk of bias and indirectness and imprecision) from 2 prospective double-blinded RCTs^{182,183} showing no benefit of seizure prophylaxis.

Post-Cardiac Arrest Seizure Treatment

For the critical outcomes of survival with favorable neurological outcome or survival at discharge/30 days or longer, we identified no RCTs or nonrandomized studies that addressed the effect of post-cardiac arrest seizure treatment, compared with no seizure treatment, on outcomes.

Treatment Recommendations

This treatment recommendation has been updated from 2015.^{1,7}

We suggest against seizure prophylaxis in adult post-cardiac arrest survivors (weak recommendation, very low-certainty evidence).

We suggest treatment of seizures in adult post-cardiac arrest survivors (weak recommendation, very low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision table is included in [Supplement Appendix A-8](#). The task force decision to suggest against post-cardiac arrest seizure prophylaxis was primarily based on the absence of direct evidence that prophylactic therapy with antiepileptic drugs prevents seizures or improves important outcomes in adult comatose cardiac arrest survivors. However, the task force did recognize the very low certainty of the evidence

from RCTs. The task force also considered that seizure prophylaxis in other forms of acute brain injury is not associated with improved outcomes, and that most drugs used for seizure prophylaxis can have significant side effects. Finally, the task force acknowledged that most comatose cardiac arrest survivors routinely receive sedatives such as propofol or benzodiazepines that are known to have antiepileptic effects. However, the task force identified no controlled studies that examined whether different sedation strategies or choices of sedation drugs had an impact on the incidence of post-cardiac arrest seizures.

The task force decision to suggest treatment of seizures in post-cardiac arrest survivors takes into consideration the absence of direct evidence that seizure treatment improves critical outcomes in this patient population. However, there are no published controlled clinical studies. Therefore, the task force weighed the fact that ongoing seizures have the potential to worsen brain injury, and treatment of recurrent seizures and status epilepticus constitutes “standard of care” in other patient populations. A large randomized trial is currently underway investigating the benefit of systematic antiepileptic drug therapy with the goal of suppressing all epileptiform activity on the EEG versus standard treatment of clinical seizures only in post-cardiac arrest status epilepticus. (TELSTAR trial [Treatment of Electroencephalographic Status Epilepticus After Cardiopulmonary Resuscitation], NCT02056236)

Indirect evidence from case series suggests that sedatives such as propofol are effective in suppressing both clinical convulsions and epileptiform activity on EEG in these patients.^{185–187} A recent retrospective study provides some evidence that conventional antiepileptic drugs (specifically valproate and levetiracetam) also have an effect in suppressing epileptiform activity in the EEG.¹⁸⁸ In a recent comparison of valproate, levetiracetam, and fosphenytoin for convulsive status epilepticus, the 3 drugs were equally effective but fosphenytoin caused more episodes of hypotension and need for tracheal intubation.¹⁸⁹ However, it is important to note that this study excluded post-cardiac arrest patients. On the basis of these results, the task force discussed using valproate and levetiracetam as first-line drugs in post-cardiac arrest seizure treatment.

There is no direct evidence of undesirable effects of antiepileptic drug therapy in comatose post-cardiac arrest survivors. Treatment with sedatives and conventional antiepileptic drugs in high doses has the potential to cause delayed awakening, prolonged need for mechanical ventilation, and increased critical care days. Importantly, generalized myoclonus in combination with epileptiform discharges may be manifestations of Lance-Adams syndrome, which is compatible with a good outcome.^{187,190} In such cases, overly aggressive

sedation and treatment with high doses of conventional antiepileptic drugs may confound the clinical examination and lead to overly pessimistic prognostication.

The relative benefit of continuous EEG compared with intermittent EEG monitoring was not specifically reviewed. Continuous EEG monitoring is labor intensive and likely to add significant cost to patient care. The net cost-effectiveness of this approach is controversial and may depend substantially on the setting.^{191,192} The task force also discussed the potential cost of delayed neurological prognostication and prolonged ICU care associated with active treatment of seizures because of the need to continue sedation.

Knowledge Gaps

- There is no high-certainty evidence of a positive effect of antiepileptic drugs on the outcome of post-cardiac arrest patients with seizures.
- There are no RCTs specifically designed to evaluate the impact of post-cardiac arrest seizure prophylaxis on the incidence of seizures and on neurological outcome.
- There are inadequate data about the timing, duration, dosing, and choice of antiepileptic drugs for seizure prophylaxis in comatose post-cardiac arrest patients.
- The utility of continuous EEG versus intermittent EEG monitoring in the diagnosis and treatment of seizures in comatose postcardiac arrest patients remains controversial.
- The threshold for treating epileptiform activity other than convulsive seizures (eg, generalized epileptiform discharges) is poorly defined.
- Standardized terminology for classification of epileptiform activity in the EEG of comatose post-cardiac arrest patients is increasingly used. There remains a need to develop consensus on the definition of post-cardiac arrest status epilepticus.
- The value of using volatile anesthetics to treat refractory status epilepticus on post-cardiac arrest patients is currently unknown.

Targeted Temperature Management (ALS 455, 790, 791, 802, 879: EvUp)

A comprehensive SysRev of TTM^{193,194} was conducted for the 2015 CoSTR.^{1,7} The task force chose to delay updating this SysRev until the completion and publication of the Targeted Hypothermia Versus Targeted Normothermia After Out-of-Hospital Cardiac Arrest (TTM2) RCT (NCT02908308). EvUps for use of TTM and TTM duration were completed and appear in [Supplement Appendix C-14 and C-15](#).

The results of the HYPERION trial (Therapeutic Hypothermia After Cardiac Arrest in Nonshockable Rhythm) were recently published.¹⁹⁵ In this French trial, 581 adult

patients who were comatose after resuscitation from either an IHCA or OHCA with an initial nonshockable rhythm were randomized to either TTM with a target temperature of 33°C or TTM with a temperature of 37°C, both for 24 hours. The primary outcome (the proportion of patients with a CPC of either 1 or 2 at 90 days after the cardiac arrest) significantly favored the 33°C group. At 90 days, 29 of 284 patients (10.2%) in the 33°C group were alive with a CPC of 1 or 2, as compared with 17 of 297 (5.7%) in the normothermia group (risk difference, 4.5%; 95% CI, 0.1–8.9; $P=0.04$). There was no difference in mortality at 90 days (81.3% versus 83.2%; risk difference, –1.9%; 95% CI, –8.0 to 4.3).

This trial does not lead to any immediate changes to the 2015 ILCOR treatment recommendations^{1,7} but reinforces the suggestion to consider TTM, targeting a constant temperature between 32°C and 36°C, in patients who remain comatose after resuscitation from either IHCA or OHCA with an initial nonshockable rhythm.

Treatment Recommendations

These treatment recommendations are unchanged from 2015.^{1,7}

We recommend selecting and maintaining a constant target temperature between 32°C and 36°C for those patients in whom temperature control is used (strong recommendation, moderate-quality evidence). Whether certain subpopulations of cardiac arrest patients may benefit from lower (32°C–34°C) or higher (36°C) temperatures remains unknown, and further research may help elucidate this.

We recommend TTM as opposed to no TTM for adults with OHCA with an initial shockable rhythm who remain unresponsive after ROSC (strong recommendation, low-quality evidence).

We suggest TTM as opposed to no TTM for adults with OHCA with an initial nonshockable rhythm who remain unresponsive after ROSC (weak recommendation, very low-quality evidence).

We suggest TTM as opposed to no TTM for adults with IHCA with any initial rhythm who remain unresponsive after ROSC (weak recommendation, very low-quality evidence).

We suggest that if TTM is used, duration should be at least 24 hours (weak recommendation, very low-quality evidence).

We recommend against routine use of prehospital cooling with rapid infusion of large volumes of cold IV fluid immediately after ROSC (strong recommendation, moderate-quality evidence).

We suggest prevention and treatment of fever in persistently comatose adults after completion of TTM between 32°C and 36°C (weak recommendation, very low-quality evidence).

Prognostication in Comatose Patients After Resuscitation From Cardiac Arrest

Combined Prognostic Systematic Reviews

Many comatose post-cardiac arrest patients will not survive or will survive with an unfavorable neurological outcome. In some regions, family and treating teams may limit or withdraw life-sustaining treatment when unfavorable neurological outcomes are expected. Therefore, reliable strategies for timely prognostication are a critical component of any cardiac arrest system of care. The 2015 CoSTR distinguished between studies of prognostication among patients treated with or without hypothermia. For this 2020 CoSTR for ALS, these treatment recommendations apply regardless of the TTM strategy used. The reason for this is that in all of the studies we assessed, the population included a mix of TTM-treated and non-TTM-treated patients, and the potential impact of TTM on prognostication could not be assessed separately.

On May 31, 2013, a new search was launched, using the search strategies used for previous SysRevs on neuroprognostication. For the SysRev informing the 2020 CoSTRs, the search included studies published from January 1, 2013, to December 31, 2019 [PROSPERO Registration: CRD42019141169].

This review identified clinical signs, neurophysiological measurements, blood biomarkers, and imaging studies that had high specificity for poor neurological outcome, defined as CPC score of 3 to 5 or mRS score of 4 to 6 at hospital discharge, 1 month, or later.

The decision to limit treatment of comatose post-cardiac arrest patients should never rely on a single prognostication element. The consensus of the task force was that in patients who remain comatose in the absence of confounders (eg, sedative drugs), a multimodal approach should be used, with all supplementary tests considered in the context of the clinical examination. The most reliable combination and timing for each assessment are still to be determined and require further research.

The SysRevs supporting this CoSTR defined prediction as imprecise when the upper limit of 95% CIs for false-positive rate was above 5%.¹⁹⁶ However, there is no universal consensus on what the acceptable limits for imprecision should be. In a recent survey of 640 medical providers, Steinberg et al¹⁹⁷ reported that 56% considered an acceptable false-positive rate for withdrawal of life sustaining treatment from patients who might otherwise have recovered was 0.1% or less. In addition, 59% of respondents felt that an acceptable false-positive rate threshold for continuing life-sustaining treatment in patients with unrecognized unrecoverable injury was 1% or less.

Clinical Examination for Prognostication (ALS 450, 713, 487: SysRev)

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults who are comatose after resuscitation from cardiac arrest (either in-hospital or out-of-hospital), regardless of target temperature
- Intervention: Pupillary light reflex (PLR), pupillometry, corneal reflex, myoclonus, and status myoclonus assessed within 1 week after cardiac arrest
- Comparator: None
- Outcome: Prediction of poor neurological outcome defined as CPC 3 to 5 or mRS 4 to 6 at hospital discharge, 1 month, or later
- Study design: Prognostic accuracy studies where the 2x2 contingency table (ie, the number of true/false negatives and positives for prediction of poor outcome) was reported, or where those variables could be calculated from reported data, are eligible for inclusion. Unpublished studies, reviews, case reports, case series, studies including fewer than 10 patients, letters, editorials, conference abstracts, and studies published in abstract form were excluded.
- Time frame: In 2015, an ILCOR evidence review identified 4 categories of predictors of neurological outcome after cardiac arrest, namely clinical examination, biomarkers, electrophysiology, and imaging. In the last 4 years, several studies have been published and new predictors have been identified, therefore the topic needs an update.
- The most recent search of the previous SysRevs on neuroprognostication was launched on May 31, 2013. We searched studies published from January 1, 2013, to December 31, 2019.
- PROSPERO Registration: CRD42019141169

Consensus on Science

Pupillary Reflex

The association of a bilaterally absent standard PLR, measured at various time points, with outcome was investigated in 17 observational studies.¹⁹⁸⁻²¹⁴ Although all of this evidence was rated as very low certainty, studies that evaluated the prognostic value of absent standard PLR at time points of 72 hours or more after ROSC had greater specificity (ranging from 90% to 100%) for unfavorable neurological outcome at time points from discharge to 12 months than studies that used the absence of PLR at less than 72 hours (specificity ranging from 48% to 92%). Sensitivity appeared to decrease when using a time point of 72 hours or more, but specificity was identified as the higher priority given the critical importance of avoiding false positives.

Pupillometry

Automated assessment of PLR can be made by measuring either of the following variables:

- The percent reduction in pupillary size, which is reported as qPLR, or
- The neurological pupil index (NPI), which is based on several variables such as pupillary size, percentage of constriction, constriction velocity, and latency.

Automated Pupillometry Using Percent of Pupillary Size Reduction (qPLR). In 3 observational studies using various time points,^{209,215,216} qPLR from 0% to 13% at 24 hours predicted poor neurological outcome from 3 months to 12 months with specificity ranging from 77.8% to 98.9% and sensitivity from 17% to 66% (certainty of evidence from moderate to very low). When evaluated at 48 hours, specificity ranged from 95.7% to 100% and sensitivity from 18.1% to 58.5% (certainty of evidence from low to very low). In 1 study of 234 patients²⁰⁹ qPLR=0% at 72 hours predicted poor neurological outcome at 3 months with 100% specificity and 4.9% sensitivity (moderate certainty of evidence).

Automated Pupillometry Using Multiple Variables (NPI). In 3 observational studies,^{209,217,218} NPI from 0 to 2.40 within 24 hours predicted poor neurological outcome from hospital discharge to 3 months with 100% specificity and sensitivity ranging from 22% to 43.9% (certainty of evidence from moderate to very low). For the same outcome, 1 study with 361 patients²⁰⁹ found that NPI 2 or less at 48 hours had 100% specificity and 18.8% sensitivity, and NPI 2 or less at 72 hours had 100% specificity, and 16.9% sensitivity (moderate certainty of evidence).

Corneal Reflex

Corneal reflex at various time points was investigated in 11 observational studies.^{198,200,202,204–206,210,211,213,214,219} Although all of the evidence was rated as very low certainty, studies that evaluated the prognostic value of absent corneal reflex at time points of 72 hours or more after ROSC had greater specificity (ranging from 89% to 100%) for unfavorable neurological outcome from hospital discharge to 12 months after ROSC than studies that used the absence of corneal reflex at less than 72 hours (specificity ranging from 25% to 89%). Sensitivity appeared to decrease when using a time point of 72 hours or more, but specificity was determined to be a higher priority given the critical importance of avoiding false positives.

Myoclonus

Presence of myoclonus within 96 hours after ROSC was investigated in 6 studies^{200,210,219–222} and predicted poor neurological outcome from hospital discharge to 6 months with specificity ranging from 77.8% to 100% and sensitivity ranging from 18.2% to 44.4% (very low-certainty evidence). However, definitions of myoclonus were provided in only 1 study.²²⁰

Status Myoclonus

Presence of status myoclonus within 72 hours after ROSC was investigated in 2 studies^{178,223} and predicted poor neurological outcome from hospital discharge to 6 months with specificity ranging from 99.8% to 100% and sensitivity ranging from 12.2% to 49.1% (very low-certainty evidence). The definitions of status myoclonus differed between these 2 studies.

Treatment Recommendations

We recommend that neuroprognostication always be undertaken by using a multimodal approach because no single test has sufficient specificity to eliminate false positives (strong recommendation, very low-certainty evidence).

We suggest using PLR at 72 hours or more after ROSC for predicting neurological outcome of adults who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence).

We suggest using quantitative pupillometry at 72 hours or more after ROSC for predicting neurological outcome of adults who are comatose after cardiac arrest (weak recommendation, low-certainty evidence).

We suggest using bilateral absence of corneal reflex at 72 hours or more after ROSC for predicting poor neurological outcome in adults who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence).

We suggest using presence of myoclonus or status myoclonus within 7 days after ROSC, in combination with other tests, for predicting poor neurological outcome in adults who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence). We also suggest recording EEG in the presence of myoclonic jerks to detect any associated epileptiform activity (weak recommendation, very low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

As noted in the previous CoSTR on this topic in 2015,^{1,7} the task force consensus is that a multimodal approach should be used in all cases with all supplementary tests considered in the context of the clinical examination.

The evidence-to-decision tables are included in [Supplement Appendixes A-9, 10, 11, and 12](#). For standard PLR, NPI, and corneal reflex, the suggestion to use these findings at 72 hours or more after ROSC was based both on the specificity found in different studies and on the perceived importance of eliminating confounding effects of sedatives or muscle relaxants as much as possible. Only some of the included studies specifically excluded the presence of residual sedation at the time the pupillary or corneal reflex was assessed.

For assessment of the pupillary reflex, the task force felt that NPI has the potential for being more accurate and less prone to bias and subjectivity. This benefit,

however, may be counterbalanced by the need for more equipment and specialized training to obtain the NPi.

Results of clinical examination usually cannot be concealed from the treating team. Therefore, a risk of self-fulfilling prophecy exists even when index tests that are based on clinical examination are not explicitly included in the criteria for withdrawal of life-sustaining therapy.

Although definitions of both myoclonus and status myoclonus are missing from most studies and are inconsistent in others, the presence of myoclonus is associated with poor outcome in patients who are comatose after ROSC from cardiac arrest and the finding may be useful within the context of a multimodal prognostic assessment. Myoclonus and status myoclonus are inconsistently associated with epileptiform activity on the EEG. Importantly, generalized myoclonus associated with favorable clinical features, such as a continuous or reactive EEG background or preserved brain stem reflexes, may be manifestations of Lance-Adams syndrome, which is compatible with a good outcome.^{187,190}

Knowledge Gaps

- Absence of residual effects from sedatives must be specifically assessed in studies evaluating the accuracy of predictors on the basis of clinical examination after cardiac arrest.
- The interrater agreement for the assessment of standard PLR, corneal reflex, and myoclonus/status myoclonus in patients resuscitated from cardiac arrest deserves investigation.
- The number of studies documenting pupillometry for predicting poor outcome after cardiac arrest is still low. A consistent threshold for 100% specificity has not been identified for qPLR or NPi.
- Achieving a uniform and consensus-based definition of both myoclonus and status myoclonus is necessary. The role of EEG as an additional tool to investigate the nature and the prognostic significance of myoclonus deserves investigation.
- The most reliable combination and timing for each assessment remains to be determined.
- The potential impact of TTM on prognostication remains to be determined.

Neurophysiological Tests for Prognostication (ALS 450, 713, 460: SysRev)

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults who are comatose after resuscitation from cardiac arrest in any setting (in-hospital or out-of-hospital) and regardless of target temperature
- Intervention: Electrophysiology studies assessed within 1 week after cardiac arrest
- Comparator: None

- Outcome: Prediction of unfavorable neurological outcome defined as CPC 3 to 5 or mRS 4 to 6 at hospital discharge, 1 month, or later
- Study design: Prognostic accuracy studies where the 2x2 contingency table (ie, the number of true/false negatives and positives for prediction of poor outcome) was reported, or where those variables could be calculated from reported data, are eligible for inclusion. Unpublished studies, reviews, case reports, case series, studies including fewer than 10 patients, letters, editorials, conference abstracts, and studies published in abstract form are excluded.
- Time frame: In 2015,^{1,7} an ILCOR evidence review identified 4 categories of predictors of neurological outcome after cardiac arrest, namely clinical examination, biomarkers, electrophysiology, and imaging. In the last 4 years, several studies have been published and new predictors have been identified, therefore the topic needs an update.
- The most recent search of the previous SysRevs on neuroprognostication was launched on May 31, 2013. We searched studies published from January 1, 2013, to December 31, 2019.
- PROSPERO Registration: CRD42019141169

Consensus on Science

Somatosensory Evoked Potentials

The prognostic value of somatosensory evoked potentials (SSEPs) was investigated in 14 observational studies.^{199,205,208–211,214,224–230} In 4 studies,^{205,224,228,229} bilaterally absent N20 SSEP wave within 24 hours after ROSC predicted poor neurological outcome from hospital discharge to 6 months. Specificity was 100% and sensitivity ranged from 33.3% to 57.7% (very low-certainty evidence). In 1 study,¹⁹⁹ an absent N20 wave on one side and an absent or low-voltage N20 wave on the other side within 24 hours after ROSC predicted poor neurological outcome at 6 months. Specificity was 100% and sensitivity was 49.6% (very low-certainty evidence).

In 12 studies,^{205,208–211,214,225–230} bilaterally absent N20 SSEP wave at 24 to 96 hours after ROSC predicted poor neurological outcome from hospital discharge to 6 months. Specificity ranged from 50% to 100% and sensitivity ranged from 18.2% to 69.1% (very low-certainty evidence).

Unreactive EEG

The prognostic value of an unreactive EEG was investigated in 10 observational studies.^{210,219,229,231–237} In 9 of these studies,^{210,219,229,232–237} an unreactive EEG within 72 hours after ROSC predicted poor neurological outcome from hospital discharge to 6 months. Specificity ranged from 41.7% to 100% and sensitivity ranged from 50% to 97.1% (certainty of evidence from moderate to very

low). Specificity was below 90% in most of these studies, reaching 100% in only 2 of them.

In 1 study,²³¹ an unreactive EEG at a median of 77 hours after ROSC (interquartile range [IQR], 53–102) predicted poor neurological outcome at 6 months with 70% specificity and 88.1% sensitivity (very low-certainty evidence).

Rhythmic/Periodic Discharges

The prognostic value of rhythmic/periodic discharges were investigated in 9 observational studies.^{199,210,228,231,237–241}

In 2 studies,^{199,238} rhythmic/periodic discharges within 24 hours after ROSC predicted poor neurological outcome from 3 months to 6 months. Specificity was 100% and sensitivity ranged from 2.4% to 7.9% (certainty of evidence from moderate to very low).

In 4 studies,^{210,228,238,239} rhythmic/periodic discharges within 48 hours after ROSC predicted poor neurological outcome from 3 months to 6 months. Specificity ranged from 97.2% to 100% and sensitivity ranged from 8.1% to 42.9% (certainty of evidence from moderate to very low).

In 3 studies,^{228,237,239} rhythmic/periodic discharges at 48 to 72 hours after ROSC predicted poor neurological outcome from 1 month to 6 months. Specificity ranged from 66.7% to 96.1% and sensitivity ranged from 11.4% to 50.8% (certainty of evidence from low to very low).

In 2 studies,^{231,240} rhythmic/periodic discharges at the median time of 76 to 77 hours after ROSC predicted poor neurological outcome at 6 months. Specificity ranged from 97% to 100% and sensitivity ranged from 5% to 40% (certainty of evidence from low to very low).

In 1 study,²⁴¹ rhythmic/periodic discharges within 5 days after ROSC predicted poor neurological outcome at 6 months. Specificity was 100% and sensitivity was 15.7% (moderate certainty of evidence).

Sporadic, Nonrhythmic/Periodic Discharges

The prognostic value of sporadic, nonrhythmic/periodic discharges was investigated in 5 observational studies.^{199,226,228,237,238} In 3 studies,^{199,226,238} sporadic, nonrhythmic/periodic discharges within 24 hours after ROSC predicted poor neurological outcome from 3 months to 6 months. Specificity ranged from 84.6% to 100% and sensitivity ranged from 0.5% to 7.9% (certainty of evidence from moderate to very low).

In 3 studies,^{226,228,238} sporadic, nonrhythmic/periodic discharges within 48 hours predicted poor neurological outcome from 3 months to 6 months. Specificity ranged from 95.8% to 99.5% and sensitivity ranged from 0.4% to 13.3% (certainty of evidence from moderate to very low).

In 3 studies,^{226,228,237} sporadic, nonrhythmic/periodic discharges at 48 to 72 hours predicted poor

neurological outcome from 1 month to 6 months. Specificity ranged from 88.9% to 97.3% and sensitivity ranging from 0.6% to 38.5% (certainty of evidence from low to very low).

In 1 study,²²⁶ sporadic, nonrhythmic/periodic discharges at 96 to 120 hours predicted poor neurological outcome at 6 months. Specificity ranged from 66.7% to 82.1% and sensitivity ranged from 17.6% to 21.3% (very low-certainty evidence).

Seizures

The prognostic implications of seizures were investigated in 5 observational studies.^{220,231,236–238} In 4 of these studies, seizures were recorded within 72 hours after ROSC, and in 1 study,²³¹ they were recorded at a median of 77 (53–102) hours after ROSC. In these studies, the presence of seizures predicted poor neurological outcome from hospital discharge to 6 months with 100% specificity and sensitivity ranging from 0.6% to 26.8% (certainty of evidence from moderate to very low).

The prognostic implications of status epilepticus were investigated in 6 studies.^{202,225,236,241–243} The definitions of status epilepticus were inconsistent across studies. In these studies, status epilepticus within 5 days after ROSC predicted poor neurological outcome from hospital discharge to 6 months. Specificity ranged from 82.6% to 100% and sensitivity ranged from 1.8% to 50% (certainty of evidence low to very low).

In 3 of these studies,^{202,225,236} EEG was recorded within 72 hours after ROSC and specificity was 100%. In another study,²⁴³ specificity was 100% only when status epilepticus originated from a discontinuous or burst-suppression background.

Burst Suppression

The possible prognostic value of burst suppression was investigated in 6 observational studies.^{202,220,225,231,233,240} In 2 studies,^{220,233} burst suppression within 24 hours after ROSC predicted poor neurological outcome to hospital discharge with 50% to 100% specificity and 50% to 51.5% sensitivity (certainty of evidence very low).

In 5 studies,^{202,225,231,233,240} burst suppression at 24 to 120 hours after ROSC predicted poor neurological outcome at hospital discharge to 6 months. Specificity ranged from 91.7% to 100% and sensitivity ranged from 13.9% to 55.6% (certainty of evidence from low to very low).

Definitions of burst suppression used in these studies varied when they were included at all. In 2 studies,^{231,240} the American Clinical Neurophysiology Society definition²⁴⁴ was used. In 1 study, a non-American Clinical Neurophysiology Society definition was used, while in the remaining studies, no specific definition was used.

Synchronous Burst Suppression. In 1 study,²²⁶ a synchronous burst suppression at 6 to 96 hours after ROSC

predicted poor neurological outcome at 6 months with 100% specificity and sensitivity ranging from 1.1% to 31.7% (certainty of evidence from moderate to low).

Heterogeneous Burst Suppression. In 1 study,²²⁶ heterogeneous burst suppression at 6 to 120 hours after ROSC predicted poor neurological outcome at 6 months. Specificity ranged from 90.7% to 100% and sensitivity ranged from 1.1% to 16.2% (certainty of evidence from moderate to very low).

Treatment Recommendations

We recommend that neuroprognostication always be undertaken by using a multimodal approach because no single test has sufficient specificity to eliminate false positives (strong recommendation, very low-certainty evidence).

We suggest using a bilaterally absent N20 wave of SSEP in combination with other indices to predict poor outcome in adult patients who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence).

We suggest against using the absence of EEG background reactivity alone to predict poor outcome in adult patients who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence).

We suggest using the presence of seizure activity on EEG in combination with other indices to predict poor outcome in adult patients who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence).

We suggest using burst suppression on EEG in combination with other indices to predict poor outcome in adult patients who are comatose and effects of sedation after cardiac arrest have cleared (weak recommendation, very low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision tables are included in [Supplement Appendixes A-13, 14, 15, 16, and 17](#).

In making a recommendation about use of SSEPs for prognostication, the task force considered that SSEPs have a low risk of confounding from TTM or sedation and a large size of effect (high precision). However, to limit the risk of self-fulfilling prophecy, combining evaluation of SSEPs with other indices of poor neurological outcome is prudent.

In almost all studies, we included the specificity of unreactive EEG background for predicting poor outcome, and its precision was low. In addition, both definitions of and stimuli to induce EEG reactivity were inconsistent across studies.

In most of the studies, we reported the specificity of rhythmic/periodic epileptiform activity for predicting poor outcome as 100%. Specificity was lower for sporadic epileptiform discharges.

In all studies, we included the specificity of American Clinical Neurophysiology Society-defined seizures on EEG for predicting poor outcome as 100%.²⁴⁴ This specificity was consistent throughout the first 72 hours after ROSC.

Specificity of status epilepticus for predicting poor outcome was 100% in only half of the studies we included. An additional challenge for use of studies of status epilepticus for prognostication is the inconsistency of its definitions in reported studies.

In all studies we included, the presence of burst suppression on EEG predicted poor neurological outcome with a specificity above 90%, and in most studies, the specificity was 100%. Because sedative agents can affect the EEG, the most prudent strategy is to assess burst suppression for prognostication when any effects of sedation medications have cleared.

Knowledge Gaps

- Further studies are needed to evaluate the added value of assessing SSEPs in combination with other predictors of poor neurological outcome after cardiac arrest.
- It is desirable that future studies adopt a standard definition of background EEG reactivity. An international consensus statement on EEG reactivity testing (eg, stimulus protocol) has been proposed.²⁴⁵
- It is desirable that future studies adopt a standard definition of epileptiform discharges.
- The specific predictive value of the different epileptiform subtypes, their prevalence, and their combination with background EEG deserves further investigation.
- Precision was low or very low in most studies of the association of seizures with outcome. Further studies are needed to confirm the predictive value of seizures for poor outcome after cardiac arrest.
- A standard definition of status epilepticus is urgently needed.
- It is desirable that future studies adopt a standard definition of burst suppression, such as the one included in the American Clinical Neurophysiology Society's Standardized Critical Care EEG Terminology.²⁴⁴
- The accuracy of synchronous burst suppression for prognostication (identical/highly epileptiform bursts) deserves further investigation.
- It is desirable to achieve a consensus definition of the term, "highly malignant EEG patterns"

in patients who are comatose after resuscitation from cardiac arrest.

- The potential impact of TTM on prognostication remains to be determined.

Blood Biomarkers for Prognostication (ALS 450, 713, 484: SysRev)

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults who are comatose after resuscitation from cardiac arrest in any setting (in-hospital or out-of-hospital) and regardless of target temperature
- Intervention: The use of neuron-specific enolase (NSE), S-100B, glial fibrillary acidic protein, serum tau protein, and neurofilament light chain assessed within 1 week after cardiac arrest
- Comparator: None
- Outcome: Prediction of unfavorable neurological outcome, defined as CPC 3 to 5 or mRS 4 to 6 at hospital discharge, 1 month, or later
- Study design: Prognostic accuracy studies where the 2×2 contingency table (ie, the number of true/false negatives and positives for prediction of poor outcome) was reported, or where those variables could be calculated from reported data, are eligible for inclusion. Unpublished studies, reviews, case reports, case series, studies including fewer than 10 patients, letters, editorials, conference abstracts, and studies published in abstract form were excluded.
- Time frame: In 2015, an ILCOR evidence review^{1,7} identified 4 categories of predictors of neurological outcome after cardiac arrest, namely clinical examination, biomarkers, electrophysiology, and imaging. In the last 4 years, several studies have been published and new predictors have been identified, therefore the topic needs an update.
- The most recent search of the previous SysRevs on neuroprognostication was launched on May 31, 2013. We searched studies published from January 1, 2013, to December 31, 2019.
- PROSPERO Registration: CRD42019141169

Consensus on Science

Neuron-Specific Enolase

The prognostic value of NSE was investigated in 12 observational studies.^{202,206,208,214,239,246–252} In these studies, NSE with thresholds ranging from 33 to 120 µg/L within 72 hours after ROSC predicted poor neurological outcome from hospital discharge to 6 months. Specificity ranged from 75% to 100% and sensitivity ranged from 7.8% to 83.6% (certainty of evidence from moderate to very low).

In 1 study,²⁴⁸ NSE with a threshold of 50.2 µg/L at day 4 (after ROSC) predicted poor neurological outcome at 1 month with 100% specificity and 42.1% sensitivity (moderate certainty of evidence).

S-100B

The accuracy of S-100B protein in predicting poor outcome in patients with ROSC after cardiac arrest was investigated in 3 observational studies.^{251,253,254}

In 2 studies,^{251,254} S-100B protein with threshold ranging from 3.58 to 16.6 µg/L immediately after ROSC predicted poor neurological outcome from 3 to 6 months with 100% specificity and sensitivity ranging from 2.8% to 26.9% (certainty of evidence from moderate to very low).

In 3 studies,^{251,253,254} S-100B protein with a threshold ranging from 0.193 to 2.59 µg/L at 24 hours after ROSC predicted poor neurological outcome from 3 to 6 months with 100% specificity and sensitivity ranging from 10.1% to 77.6% (certainty of evidence from moderate to very low). In the same 3 studies,^{251,253,254} S-100B protein with a threshold ranging from 0.159 to 3.67 µg/L at 48 hours after ROSC predicted poor neurological outcome from 3 to 6 months with 100% specificity and sensitivity ranging from 5% to 77.6% (certainty of evidence from moderate to very low). In the same 3 studies,^{251,253,254} S-100B protein with a threshold ranging from 0.202 to 1.83 µg/L at 72 hours after ROSC predicted poor neurological outcome from 3 to 6 months with 100% specificity and sensitivity ranging from 5% to 61.2% (certainty of evidence from moderate to very low).

Glial Fibrillary Acidic Protein

In 1 study,²⁵² glial fibrillary acidic protein with a threshold of 0.08 µg/L at 48±12 hours after ROSC predicted poor neurological outcome at 1 month with 100% specificity and 21.3% sensitivity (low-certainty evidence).

Serum Tau Protein

In 1 study with 667 patients,²⁵⁵ serum tau protein with a threshold ranging from 72.7 to 874.5 ng/L at 24 to 72 hours after ROSC predicted poor neurological outcome at 6 months with 100% specificity and sensitivity ranging from 4% to 42% (very low-certainty evidence).

Serum Neurofilament Light Chain

In 1 study,²⁵⁶ serum neurofilament light chain with a threshold ranging from 1539 to 12317 pg/mL at 24 to 72 hours after ROSC predicted poor neurological outcome at 6 months with 100% specificity and sensitivity ranging from 53.1% to 65% (moderate certainty of evidence).

In 1 study,²⁵⁷ serum neurofilament light chain with a threshold ranging from 252 to 405 pg/mL from day 1 to day 7 after ROSC predicted poor neurological outcome (CPC 4–5) at 6 months with 100% specificity and sensitivity ranging from 55.6% to 94.4% (very low-certainty evidence).

Treatment Recommendations

We recommend that neuroprognostication always be undertaken by using a multimodal approach because no single test has sufficient specificity to eliminate false positives (strong recommendation, very low-certainty evidence).

We suggest using NSE within 72 hours after ROSC, in combination with other tests, for predicting neurological outcome of adults who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence). There is no consensus on a threshold value.

We suggest against using S-100B protein for predicting neurological outcome of adults who are comatose after cardiac arrest (weak recommendation, low-certainty evidence).

We suggest against using serum levels of glial fibrillary acidic protein, serum tau protein, or neurofilament light chain for predicting poor neurological outcome of adults who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

As was noted in the information addressing this topic in the 2015 CoSTR,^{1,7} the task force opinion is that a multimodal approach should be used in all cases with all supplementary tests considered in the context of prognostication.

The evidence-to-decision tables are included in [Supplement Appendixes A-18, 19, and 20](#).

Limited evidence suggests that high concentrations of NSE predict poor neurological outcome with 100% specificity at 24 to 72 hours after cardiac arrest, but there is a wide variability of thresholds for 100% specificity across studies. Lack of blinding was a limitation in most of included studies, even if withdrawal of life sustaining therapy based only on NSE was not documented.

Although the risk of self-fulfilling prophecy for S-100B protein is lower than that observed in other predictors, the evidence is limited by the few available studies and the wide variability of thresholds for 100% specificity across studies.

The supporting evidence about the use of neurofilament light chain, glial fibrillary acidic protein, and serum tau protein for prognostication after cardiac arrest is limited to very few studies. Consistent thresholds for 100% specificity need to be identified before any of these biomarkers can be recommended for prognostication in the clinical setting. These biomarker tests are not widely available. The methods used for measuring these biomarkers need to be more widely available, standardized, and studied.

Knowledge Gaps

- Large cohort studies are desirable to identify consistent NSE and S-100B thresholds for predicting poor neurological outcome after cardiac arrest. There is very little evidence concerning the

predictive value of these biomarkers when measured later than 72 hours after ROSC.

- Further studies on glial fibrillary acidic protein, serum tau protein, and neurofilament light chain are needed to confirm their predictive value after cardiac arrest, to assess their reproducibility, and to identify consistent thresholds for 100% specificity.
- The potential impact of TTM on prognostication remains to be determined.

Imaging for Prognostication (ALS 450, 713, 458: SysRev)

Population, Intervention, Comparator, Outcome, Study Design, and Time Frame

- Population: Adults who are comatose after resuscitation from cardiac arrest in any setting (in-hospital or out-of-hospital) and regardless of target temperature
- Intervention: Imaging studies assessed within 1 week after cardiac arrest
- Comparator: None
- Outcome: Unfavorable neurological outcome defined as CPC 3 to 5 or mRS 4 to 6 at hospital discharge, 1 month, or later
- Study design: Prognostic accuracy studies where the 2x2 contingency table (ie, the number of true/false negatives and positives for prediction of poor outcome) was reported, or where those variables could be calculated from reported data, are eligible for inclusion. Unpublished studies, reviews, case reports, case series, studies including fewer than 10 patients, letters, editorials, conference abstracts, and studies published in abstract form were excluded.
- Time frame: In 2015,^{1,7} an ILCOR evidence review identified 4 categories of predictors of neurological outcome after cardiac arrest, namely clinical examination, biomarkers, electrophysiology, and imaging. In the last 4 years, several studies have been published and new predictors have been identified, therefore the topic needs an update.
- The most recent search of the previous SysRevs on neuroprognostication was launched on May 31, 2013. We searched studies published from January 1, 2013, to December 31, 2019.
- PROSPERO Registration: CRD42019141169

Consensus on Science

Gray Matter-to-White Matter Ratio

Gray Matter-to-White Matter Ratio: Average. The prognostic value of the gray matter-to-white matter ratio (GWR) average was investigated in 7 observational studies.^{203,214,258–262} In 4 studies,^{214,260,261,263} a GWR

Table 3a. Sensitivity and Specificity of GWR at 50 (IQR, 26–107) Minutes From ROSC by Brain Location in Patients With Cardiac Arrest of Cardiac Etiology

Study, Year	GWR	Location or Type	Sensitivity	Specificity	Certainty of Evidence
Poor Neurological Outcome at Discharge					
Lee, 2015 ²⁶⁶	≤1.13	Average	3.5%	100%	Very low
	≤1.11	Basal ganglia	3.5%	100%	Very low
	≤1.107	Putamen/corpus callosum	5.6%	100%	Very low
	≤1.06	Simplified	3.5%	100%	Very low
	≤1.094	Caudate nucleus/posterior limb of the internal capsule	3.5%	100%	Very low
	≤1.15	Cerebrum	4.2%	100%	Very low

GWR indicates gray matter-to-white matter ratio; and IQR, interquartile range.

average 1.23 or less within 6 hours after ROSC predicted poor neurological outcome from hospital discharge to 6 months with 100% specificity and sensitivity ranging from 13.3% to 83.8% (certainty of evidence from low to very low).

In 1 study,²⁰³ a GWR average 1.13 or less at 124.5±59.9 minutes from ROSC predicted poor neurological outcome at 1 month with 85% specificity and 29.8% sensitivity (very low-certainty evidence).

In 1 study,²⁵⁹ a GWR average 1.077 or less within 24 hours after ROSC predicted poor neurological outcome at hospital discharge with 100% specificity and 15.6% sensitivity (very low-certainty evidence).

In 1 study,²⁵⁸ a GWR average 1.14 or less within 72 hours after ROSC predicted poor neurological outcome at hospital discharge with 100% specificity and 38.1% sensitivity (very low-certainty evidence).

Gray Matter-to-White Matter Ratio: Basal Ganglia. The prognostic value of the GWR in the basal ganglia was investigated in 4 observational studies.^{199,258,261,264} In 1 study,²⁶¹ GWR-basal ganglia 1.12 or less within 1 hour after ROSC predicted poor neurological outcome at hospital discharge with 100% specificity and 3.3% sensitivity (very low-certainty evidence).

In 2 studies,^{199,264} GWR-basal ganglia 1.21 or less within 24 hours after ROSC predicted poor neurological

outcome at 6 months with 100% specificity and sensitivity ranging from 41.8% to 42.1% (certainty of evidence from moderate to very low).

In 1 study,²⁵⁸ GWR-basal ganglia 1.12 or less within 72 hours after ROSC predicted poor neurological outcome at hospital discharge with 100% specificity and 28.6% sensitivity (very low-certainty evidence).

Gray Matter-to-White Matter Ratio: Putamen/Corpus Callosum. The prognostic value of the GWR putamen/corpus callosum was investigated in 3 observational studies.^{247,260,265}

In 2 studies,^{247,260} the GWR putamen/corpus callosum 1.17 or less within 6 hours after ROSC predicted poor neurological outcome from hospital discharge to 6 months with 100% specificity and sensitivity ranging from 31.3% to 52.9% (very low-certainty evidence).

In 1 study,²⁶⁵ of 258 patients, the GWR putamen/corpus callosum 0.91 or less within 24 hours after ROSC predicted poor neurological outcome at 6 months with 100% specificity and 1.7% sensitivity (very low-certainty evidence).

Gray Matter-to-White Matter Ratio: Simplified (Putamen/Posterior Limb of Internal Capsule). In 1 observational study, GWR-simplified²⁵⁸ a ratio 1.1 or less

Table 3b. Sensitivity and Specificity of GWR at 67 (IQR, 29–115) Minutes From ROSC by Brain Location in Patients With Cardiac Arrest of Noncardiac Etiology

Study	GWR	Location or Type	Sensitivity	Specificity	Certainty of Evidence
Poor Neurological Outcome at Discharge					
Lee, 2016 ²⁶⁷	≤1.22	Average	28.3%	100%	Very low
	≤1.17	Basal ganglia	26.2%	100%	Very low
	≤1.2	Putamen/corpus callosum	43.4%	100%	Very low
	≤1.12	Simplified	9.7%	100%	Very low
	≤1.138	Caudate nucleus/posterior limb of the internal capsule	20%	100%	Very low
	≤1.2	Cerebrum	11%	100%	Very low

GWR indicates gray matter-to-white matter ratio; and IQR, interquartile range.

within 72 hours after ROSC predicted poor neurological outcome at hospital discharge with 100% specificity and 28.6% sensitivity (very low-certainty evidence).

Gray Matter-to-White Matter Ratio: Caudate Nucleus/Posterior Limb of Internal Capsule. In 2 observational studies,^{247,260} a GWR in the caudate nucleus/posterior limb of the internal capsule 1.15 or less within 6 hours after ROSC predicted poor neurological outcome from hospital discharge to 6 months with 100% specificity and sensitivity ranging from 19.8% to 40.6% (very low-certainty evidence).

Gray Matter-to-White Matter Ratio: Cerebrum. The prognostic value of the GWR in the cerebrum was investigated in 2 observational studies.^{258,261} In 1 study,²⁶¹ a GWR in the cerebrum 1.12 or less within 1 hour after ROSC predicted poor neurological outcome at hospital discharge with 100% specificity and 20% sensitivity (very low-certainty evidence).

In 1 study,²⁵⁸ a GWR in the cerebrum 1.09 or less within 72 hours after ROSC predicted poor neurological outcome at hospital discharge with 100% specificity and 28.6% sensitivity (very low-certainty evidence).

Gray Matter-to-White Matter Ratio: Thalamus/Corpus Callosum. In 1 observational study,²⁶⁰ a GWR in the cerebrum thalamus/corpus callosum 1.13 or less at a median time of 90 (IQR, 52–150) minutes after ROSC predicted poor neurological outcome at 6 months with 100% specificity and 50% sensitivity (very low-certainty evidence).

Gray Matter-to-White Matter Ratio: Caudate Nucleus/Corpus Callosum. In 1 observational study,²⁶⁰ the GWR in the caudate nucleus/corpus callosum 1.15 or less at median time of 90 (IQR, 52–150) minutes after ROSC predicted poor neurological outcome at 6 months with 100% specificity and 46.9% sensitivity (very low-certainty evidence).

Gray Matter-to-White Matter Ratio in Cardiac Versus Noncardiac Etiology. One study assessed the predictive value of GWR specifically in patients with cardiac arrest of cardiac etiology, and one other focused exclusively on cardiac arrest with noncardiac etiology.^{266,267} Both of these studies reported GWRs that had 100% specificity for poor neurological outcome, and sensitivity was low in all cases. Results are presented in detail in Tables 3a and 3b.

Diffusion-Weighted MRI

The prognostic value of diffusion-weighted magnetic resonance imaging (MRI) was investigated in 5 observational studies.^{198,214,260,268,269}

In 1 study,²⁶⁰ high signal intensity on diffusion-weighted MRI within 6 hours after ROSC predicted poor neurological outcome at 6 months with 100% specificity and 81.3% sensitivity (very low-certainty evidence).

In 4 studies,^{198,214,268,269} positive findings on diffusion-weighted MRI within 5 days after ROSC predicted poor neurological outcome from hospital discharge to 6 months with specificity ranging from 55.7% to 100% and sensitivity ranging from 26.9% to 92.6% (very low-certainty evidence).

Apparent Diffusion Coefficient

The prognostic value of apparent diffusion coefficient (ADC) was investigated in 2 studies.^{261,269a}

In 1 study,²⁷⁰ a mean ADC 726×10^{-6} mm²/s or less at less than 48 hours after ROSC predicted poor neurological outcome at 6 months with 100% specificity and 44% sensitivity (very low-certainty evidence).

In the same study,²⁷⁰ a mean ADC 627×10^{-6} mm²/s or less at 48 hours to 7 days after ROSC predicted poor neurological outcome at 6 months with 100% specificity and 20.8% sensitivity (very low-certainty evidence).

In the same study,²⁷⁰ an ADC volume proportion (400×10^{-6} mm²/s) greater than 2.5% at less than 48 hours after ROSC predicted poor neurological outcome at 6 months with 100% specificity and 64% sensitivity (very low-certainty evidence).

In the same study,²⁷⁰ an ADC volume proportion (400×10^{-6} mm²/s) greater than 1.66% at 48 hours to 7 days after ROSC predicted poor neurological outcome at 6 months with 100% specificity and 79.2% sensitivity (very low-certainty evidence).

In another study,^{269a} maximum cluster size in different cerebral regions on MRI 151.7×10^{-6} mm²/s or less at 46 (IQR, 37–52) hours after ROSC predicted poor neurological outcome at 6 months with 100% specificity and sensitivity ranging from 62.5% to 90% (very low-certainty evidence).

In that same study,^{269a} the lowest mean ADC in different cerebral regions on MRI 555.7×10^{-6} mm²/s or less at 46 (IQR, 37–52) hours after ROSC predicted poor neurological outcome at 6 months with 100% specificity and sensitivity ranging from 50% to 72.5% (very low-certainty evidence).

In the same study,^{269a} the lowest minimum ADC in different cerebral regions MRI 466.8×10^{-6} mm²/s or less at 46 (IQR, 37–52) hours after ROSC predicted poor neurological outcome at 6 months with 100% specificity and sensitivity ranging from 42.5% to 82.5% (very low-certainty evidence).

Treatment Recommendations

We recommend that neuroprognostication always be undertaken by using a multimodal approach because no single test has sufficient specificity to eliminate false positives (strong recommendation, very low-certainty evidence).

We suggest using GWR on brain computed tomography for predicting neurological outcome of adults who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence).

However, no GWR threshold for 100% specificity can be recommended.

We suggest using diffusion-weighted brain MRI for predicting neurological outcome of adults who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence).

We suggest using ADC on brain MRI for predicting neurological outcome of adults who are comatose after cardiac arrest (weak recommendation, very low-certainty evidence).

Justification and Evidence-to-Decision Framework Highlights

The evidence-to-decision tables are included [Supplement Appendixes A-21, 22, and 23](#). As noted in the 2015 CoSTR on this topic,^{1,7} the task force consensus is that a multimodal approach should be used in all cases with all supplementary tests considered in the context of prognostication.

In patients who are comatose after cardiac arrest, severe brain edema predicts poor outcome with high specificity. Calculation of GWR allows a quantitative evaluation of brain edema. However, there is a wide heterogeneity of measurement techniques (sites and calculation methods) for GWR. This may partly explain the wide variability of thresholds for 100% specificity across the identified studies. The evidence supporting use of the GWR for prognostication has very low certainty.

Assessing diffusion-weighted imaging has potential for predicting poor neurological outcome after cardiac arrest. The definition of a positive diffusion weighted magnetic resonance image after cardiac arrest was inconsistent or even absent in the identified studies. The supporting evidence had very low certainty.

Assessing ADC has a potential for predicting poor neurological outcome after cardiac arrest with high sensitivity. There is a wide heterogeneity of measurement techniques (sites and calculation methods) for ADC across studies. The supporting evidence for ADC had very low certainty.

Knowledge Gaps

- A consistent GWR threshold for predicting poor neurological outcome after cardiac arrest should be identified.
- A standardization of the methods for GWR calculation is warranted.
- The optimal timing for prognostication using brain computed tomography after cardiac arrest is still unknown. Studies assessing serial brain computed tomography after cardiac arrest are desirable.
- The criteria for defining a positive diffusion-weighted MRI after cardiac arrest need to be standardized.

- A consistent ADC threshold for predicting poor neurological outcome after cardiac arrest should be identified.
- Standardization of the methods for ADC calculation is needed.
- The potential impact of TTM on prognostication remains to be determined.

ALS COSTR TOPICS NOT REVIEWED IN 2020

Post-ROSC Percutaneous Coronary Intervention

Updates to 2015 CoSTRs for acute coronary syndromes (ACS) are now part of ALS postresuscitation care because there is no longer an ACS Task Force.^{271,272} The topics of percutaneous coronary intervention after ROSC in patients with and without ST-segment elevation (ACS 340, ACS 885) will be addressed in the 2021 CoSTR after publication of an ongoing SysRev.

Organ Donation After Cardiac Arrest

The 2015 treatment recommendations^{1,7} have not been updated for 2020. An ILCOR scientific statement on organ donation after OHCA will provide a narrative summary of the world literature on the incidence and outcomes of organ donation after OHCA as well as an estimation of potential donors and published implementation strategies with or without extracorporeal resuscitation. The statement includes a review of the international ethical issues and provides cost effectiveness estimates. It will make summary suggestions for implementation as well as identify key knowledge gaps that need to be addressed by future research.

Manual Defibrillation Topics Not Reviewed in 2020

- Algorithm for transition from shockable to non-shockable rhythm and vice versa (ALS 444)
- Biphasic waveforms (ALS 470)
- Pulsed biphasic waveforms (ALS 470)
- First shock energy (ALS 470)
- Single shocks versus stacked shocks (ALS 470)
- Fixed versus escalating defibrillation energy (ALS 470)
- Cardioversion strategies with implantable cardioverter-defibrillators or pacemakers (ALS 475)

Circulatory Support Topics Not Reviewed in 2020

- IABP versus manual CPR (ALS 724)
- Open-chest CPR (ALS 574)

- Impedance threshold device (ALS 579)
- Mechanical CPR devices (ALS 782)

Drugs During CPR Topics Not Reviewed in 2020

- IV fluids during cardiac arrest (ALS 578)
- Drugs for atrial fibrillation (ALS 466)
- Drugs for narrow complex tachycardia (ALS 463)
- Drugs for monomorphic wide complex tachycardia (ALS 464)
- Drugs for undifferentiated stable wide complex tachycardia (ALS 583)
- Drugs for bradycardia (ALS 465)
- Atropine for cardiac arrest (ALS 491)
- Calcium for cardiac arrest (ALS 482)

Intra-arrest Monitoring Topics Not Reviewed in 2020

- Point-of-care echocardiography for diagnosis during CPR (ALS 658)

Special Circumstances Topics Not Reviewed in 2020

- Cardiac tamponade (ALS 478)
- Cardiac arrest during coronary catheterization (ALS 479)
- Cardiac arrest in operating room (ALS 812)
- Post-op cardiothoracic surgery cardiac arrest (ALS 572)
- Electrolyte disturbances (ALS 456)
- Digoxin toxicity (ALS 468)
- Tricyclic antidepressant toxicity (ALS 429)
- Cyanide toxicity (ALS 471)
- Cocaine toxicity (ALS 474)
- Carbon monoxide toxicity (ALS 480)
- Calcium channel blocker toxicity (ALS 481)
- Beta blocker toxicity (ALS 485)
- Benzodiazepine toxicity (ALS 486)
- Lipid therapy for cardiac arrest secondary to drug toxicity (ALS 834)
- Avalanche victims (ALS 489)
- Morbid obesity (ALS 452)
- Asthma and cardiac arrest (ALS 492)
- Cardiac arrest caused by anaphylaxis (ALS 494)

Postresuscitation Care Topics Not Reviewed in 2020

- IV fluids after cardiac arrest (ALS 577)
- Mechanical circulatory support postresuscitation (ALS 447)
- Glucose control after resuscitation (ALS 580)
- Hemofiltration postresuscitation (ALS 453)
- Percutaneous coronary intervention after ROSC with ST-segment elevation (ACS 340)
- Percutaneous coronary intervention after ROSC without ST-segment elevation (ACS 885)
- Organ donation (ALS 449)

ARTICLE INFORMATION

The American Heart Association requests that this document be cited as follows: Berg KM, Soar J, Andersen LW, Böttiger BW, Cacciola S, Callaway CW, Couper K, Cronberg T, D'Arrigo S, Deakin CD, Donnino MW, Drennan IR, Granfeldt A, Hoedemaekers CWE, Holmberg MJ, Hsu CH, Kamps M, Musiol S, Nation K, Neumar RW, Nicholson T, O'Neil BJ, Otto Q, Paiva EF, Parr MJA, Reynolds JC, Sandroni C, Scholefield BR, Skrifvars MB, Wang T-L, Wetsch WA, Yeung J, Morley PT, Morrison LJ, Welsford M, Hazinski MF, Nolan JP; on behalf of the Adult Advanced Life Support Collaborators. Adult advanced life support: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2020;142(suppl 1):S92–S139. doi: 10.1161/CIR.0000000000000893

Supplemental materials are available with this article at <https://www.ahajournals.org/doi/suppl/10.1161/CIR.0000000000000893>

This article has been copublished in *Resuscitation*. Published by Elsevier Ireland Ltd. All rights reserved.

Authors

Katherine M. Berg, MD; Jasmeet Soar, MA, MB, BChir; Lars W. Andersen, MD, MPH, PhD, DMSc; Bernd W. Böttiger, MD, ML, DEAA; Sofia Cacciola, MD; Clifton W. Callaway, MD, PhD; Keith Couper, RN, PhD; Tobias Cronberg MD, PhD; Sonia D'Arrigo MD, PhD; Charles D. Deakin, MD; Michael W. Donnino, MD; Ian R. Drennan, ACP, BScHK, PhD(c); Asger Granfeldt, MD, PhD, DMSc; Cornelia W.E. Hoedemaekers MD, PhD; Mathias J. Holmberg, MD, MPH; Cindy H. Hsu MD, PhD; Marlijn Kamps, MD; Szymon Musiol, MA, MB, BChir, MRCP(UK); Kevin J. Nation, NZRN; Robert W. Neumar, MD, PhD; Tonia Nicholson, MBBS, BScPsych; Brian J. O'Neil, MD; Quentin Otto, MB, BChir; Edison Ferreira de Paiva, MD, PhD; Michael J.A. Parr, MB, BS; Joshua C. Reynolds, MD, MS; Claudio Sandroni, MD; Barnaby R. Scholefield, MBBS, MRCPCH, PhD; Markos B. Skrifvars, MD, PhD; Tzong-Luen Wang, MD, PhD, JM; Wolfgang A. Wetsch, MD; Joyce Yeung, MBChB, PhD; Peter T. Morley, MBBS; Laurie J. Morrison, MD, MSC; Michelle Welsford, BSc, MD; Mary Fran Hazinski, RN, MSN; Jerry P. Nolan, MBChB; on behalf of the Adult Advanced Life Support Collaborators

Acknowledgments

The authors thank the following individuals (the Adult Advanced Life Support Collaborators) for their contributions: Mahmoud Issa, Monica E. Kleinman, Giuseppe Ristagno, Julie Arafah, Justin L. Benoit, Maureen Chase, Bryan L. Fishberg, Gustavo E. Flores, Mark S. Link, Joseph P. Ornato, Sarah M. Perman, Comilla Sasson, and Carolyn M. Zelop.

Disclosures**Writing Group Disclosures**

Writing Group Member	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Expert Witness	Ownership Interest	Consultant/Advisory Board	Other
Katherine M. Berg	Beth Israel Deaconess Medical Center	NHLBI Grant K23 HL128814†	None	None	None	None	None	None
Jasmeet Soar	Southmead Hospital North Bristol NHS Trust (United Kingdom)	None	None	None	None	None	None	None
Lars W. Andersen	Aarhus University (Denmark)	None	None	None	None	None	None	None
Bernd W. Böttiger	University Hospital of Cologne (Germany)	None	None	FomF*; Zoll*; C.R. Bard*	None	None	None	None
Sofia Cacciola	Self-employed	None	None	None	None	None	None	None
Clifton W. Callaway	University of Pittsburgh	NIH†	None	None	None	None	None	None
Keith Couper	Warwick Medical School, University of Warwick (United Kingdom)	None	None	None	None	None	None	None
Tobias Cronberg	Lund University, Skane University Hospital (Sweden)	None	None	None	None	None	None	None
Sonia D'Arrigo	Catholic University (Italy)	None	None	None	None	None	None	None
Charles D. Deakin	NIHR Southampton Respiratory Biomedical Research Unit	None	None	None	None	None	None	None
Michael W. Donnino	Beth Israel Deaconess Medical Center	NIH†; General Electric (Investigator initiated study with VO2)*; Kaneka (Investigator initiated research grant with ubiquinol)*	None	None	None	None	None	None
Ian R. Drennan	Sunnybrook Health Sciences Center (Canada)	None	None	None	None	None	None	None
Asger Granfeldt	Aarhus University (Denmark)	None	None	None	None	None	None	None
Mary Fran Hazinski	Vanderbilt University	None	None	None	None	None	American Heart Association†	None
Cornelia W.E. Hoedemaekers	RadboudUMC (The Netherlands)	None	None	None	None	None	None	None
Mathias Johan Holmberg	Aarhus University (Denmark)	None	None	None	None	None	None	None
Cindy H. Hsu	University of Michigan	NIH (K12HL133304-01)*; AHA (AHA Strategic Network - Michigan Resuscitation Innovation and Science Enterprise (M-RISE))*	None	None	None	None	None	None
Marlijn Kamps	Radboud University (The Netherlands)	None	None	None	None	None	None	None
Peter T. Morley	University of Melbourne, Royal Melbourne Hospital (Australia)	None	None	None	None	None	None	None
Laurie J. Morrison	St Michael's Hospital, University of Toronto (Canada)	None	None	None	None	None	None	None
Szymon Musiol	Core Medical Trainee, Severn Deanery, Bristol (United Kingdom)	None	None	None	None	None	None	None

(Continued)

Writing Group Disclosures Continued

Writing Group Member	Employment	Research Grant	Other Research Support	Speakers' Bureau/ Honoraria	Expert Witness	Ownership Interest	Consultant/ Advisory Board	Other
Kevin J. Nation	New Zealand Resuscitation Council (New Zealand)	None	None	None	None	None	None	None
Robert W. Neumar	University of Michigan	NIH (R01 HL133129, K12 HL133304)†; AHA (19SFRN34760762)†	Stryker/ PhysioControl (Equipment support for laboratory and clinical research)*	None	None	None	None	None
Tonia Nicholson	Waikato Hospital Emergency Medicine (New Zealand)	None	None	None	None	None	None	None
Jerry P. Nolan	Warwick Medical School, University of Warwick (United Kingdom)	None	None	None	None	None	None	None
Brian J. O'Neil	Wayne State University	SIREN Network (Clinical trial network through NHLBI)*	None	Zoll circulation*; Genentech*	None	None	None	None
Quentin Otto	Anaesthetic Trainee, Severn Deanery, Bristol (United Kingdom)	None	None	None	None	None	None	None
Edison Ferreira de Paiva	University of Sao Paulo (Brazil)	None	None	None	None	None	None	None
Michael J.A. Parr	Liverpool Hospital, University of New South Wales and Macquarie University Hospital, Macquarie University (Australia)	None	None	None	None	None	None	None
Joshua C. Reynolds	Michigan State University	None	None	None	None	None	None	None
Claudio Sandroni	Università Cattolica del Sacro Cuore, Policlinico Gemelli (Italy)	None	None	None	None	None	None	None
Barnaby R. Scholefield	University of Birmingham (United Kingdom)	NIHR UK (Clinician Scientist Fellowship program for research into neuroprognostication after paediatric cardiac arrest)†	None	None	None	None	None	Salary (NIHR UK Fellowship Funding—Clinician Scientist (academic, non-commercial))†
Markus B. Skrifvars	Helsinki University Hospital and University of Helsinki (Finland)	COVIDIEN (Invos) (Support for study conducted in 2016–2017)*	None	BARD Medical (Ireland)*	None	None	None	None
Tzong-Luen Wang	Chang Bing Show Chwang Memorial Hospital (Taiwan)	None	None	None	None	None	None	None
Michelle Welsford	Center for Paramedic Education and Research (Canada)	None	None	None	None	None	None	None
Wolfgang A. Wetsch	University Hospital of Cologne (Germany)	None	None	None	None	None	None	None
Joyce Yeung	University of Warwick, Warwick Medical School (United Kingdom)	None	None	None	None	None	None	None

This table represents the relationships of writing group members that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all members of the writing group are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$10 000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$10 000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

*Modest.

†Significant.

Reviewer Disclosures

Reviewer	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Expert Witness	Ownership Interest	Consultant/Advisory Board	Other
Alix Carter	Dalhousie University (Canada)	Maritime Heart*	None	None	None	None	None	None
Henry Halperin	Johns Hopkins University	Zoll Medical†; NIH†	None	None	None	None	None	None
Jonathan Jui	Oregon Health and Science University	None	None	None	None	None	None	None
Fred Severyn	Denver Health and Hospital Authority and University of Colorado Campus; University of Arkansas	None	None	None	None	None	None	None
Robert A. Swor	William Beaumont Hospital	None	None	None	None	None	None	None
Andrew H. Travers	Emergency Health Services, Nova Scotia (Canada)	None	None	None	None	None	None	None

This table represents the relationships of reviewers that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all reviewers are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$10 000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$10 000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

*Modest.

†Significant.

REFERENCES

- Soar J, Callaway CW, Aibiki M, Böttiger BW, Brooks SC, Deakin CD, Donnino MW, Drajer S, Kloeck W, Morley PT, et al; on behalf of the Advanced Life Support Chapter Collaborators. Part 4: advanced life support: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2015;95:e71–e120. doi: 10.1016/j.resuscitation.2015.07.042
- Soar J, Maconochie I, Wyckoff MH, Olasveengen TM, Singletary EM, Greif R, Aickin R, Bhanji F, Donnino MW, Mancini ME, Wyllie JP, Zideman D, Andersen LW, Atkins DL, Aziz K, Bendall J, Berg KM, Berry DC, Bigham BL, Bingham R, Couto TB, Böttiger BW, Borra V, Bray JE, Breckwoldt J, Brooks SC, Buick J, Callaway CW, Carlson JN, Cassan P, Castrén M, Chang WT, Charlton NP, Cheng A, Chung SP, Considine J, Couper K, Dainty KN, Dawson JA, de Almeida MF, de Caen AR, Deakin CD, Drennan IR, Duff JP, Epstein JL, Escalante R, Gazmuri RJ, Gilfoyle E, Granfeldt A, Guerguerian AM, Guinsburg R, Hatanaka T, Holmberg MJ, Hood N, Hosono S, Hsieh MJ, Isayama T, Iwami T, Jensen JL, Kapadia V, Kim HS, Kleinman ME, Kudenchuk PJ, Lang E, Lavonas E, Liley H, Lim SH, Lockey A, Lofgren B, Ma MH, Markenson D, Meaney PA, Meyran D, Mildenhall L, Monsieurs KG, Montgomery W, Morley PT, Morrison LJ, Nadkarni VM, Nation K, Neumar RW, Ng KC, Nicholson T, Nikolaou N, Nishiyama C, Nuthall G, Ohshimo S, Okamoto D, O'Neil B, Ong GY, Paiva EF, Parr M, Pellegrino JL, Perkins GD, Perlman J, Rabi Y, Reis A, Reynolds JC, Ristagno G, Roehr CC, Sakamoto T, Sandroni C, Schexnayder SM, Scholefield BR, Shimizu N, Skrifvars M, Smyth MA, Stanton D, Swain J, Szyld E, Tijssen J, Travers A, Trevisanuto D, Vaillancourt C, Van de Voorde P, Velaphi S, Wang TL, Weiner G, Welsford M, Woodin JA, Yeung J, Nolan JP, Hazinski MF. 2019 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations Summary. *Resuscitation*. 2019;145:95–150. doi: 10.1016/j.resuscitation.2019.10.016
- Soar J, Maconochie I, Wyckoff MH, Olasveengen TM, Singletary EM, Greif R, Aickin R, Bhanji F, Donnino MW, Mancini ME, Wyllie JP, Zideman D, Andersen LW, Atkins DL, Aziz K, Bendall J, Berg KM, Berry DC, Bigham BL, Bingham R, Couto TB, Böttiger BW, Borra V, Bray JE, Breckwoldt J, Brooks SC, Buick J, Callaway CW, Carlson JN, Cassan P, Castrén M, Chang WT, Charlton NP, Cheng A, Chung SP, Considine J, Couper K, Dainty KN, Dawson JA, de Almeida MF, de Caen AR, Deakin CD, Drennan IR, Duff JP, Epstein JL, Escalante R, Gazmuri RJ, Gilfoyle E, Granfeldt A, Guerguerian AM, Guinsburg R, Hatanaka T, Holmberg MJ, Hood N, Hosono S, Hsieh MJ, Isayama T, Iwami T, Jensen JL, Kapadia V, Kim HS, Kleinman ME, Kudenchuk PJ, Lang E, Lavonas E, Liley H, Lim SH, Lockey A, Lofgren B, Ma MH, Markenson D, Meaney PA, Meyran D, Mildenhall L, Monsieurs KG, Montgomery W, Morley PT, Morrison LJ, Nadkarni VM, Nation K, Neumar RW, Ng KC, Nicholson T, Nikolaou N, Nishiyama C, Nuthall G, Ohshimo S, Okamoto D, O'Neil B, Yong-Kwang Ong G, Paiva EF, Parr M, Pellegrino JL, Perkins GD, Perlman J, Rabi Y, Reis A, Reynolds JC, Ristagno G, Roehr CC, Sakamoto T, Sandroni C, Schexnayder SM, Scholefield BR, Shimizu N, Skrifvars M, Smyth MA, Stanton D, Swain J, Szyld E, Tijssen J, Travers A, Trevisanuto D, Vaillancourt C, Van de Voorde P, Velaphi S, Wang TL, Weiner G, Welsford M, Woodin JA, Yeung J, Nolan JP, Hazinski MF. 2019 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations Summary. *Resuscitation*. 2019;145:95–150. doi: 10.1016/j.resuscitation.2019.10.016
- Soar J, Donnino MW, Maconochie I, Aickin R, Atkins DL, Andersen LW, Berg KM, Bingham R, Böttiger BW, Callaway CW, Couper K, Couto TB, de Caen AR, Deakin CD, Drennan IR, Guerguerian AM, Lavonas EJ, Meaney PA, Nadkarni VM, Neumar RW, Ng KC, Nicholson TC, Nuthall GA, Ohshimo S, O'Neil BJ, Ong GY, Paiva EF, Parr MJ, Reis AG, Reynolds JC, Ristagno G, Sandroni C, Schexnayder SM, Scholefield BR, Shimizu N, Tijssen JA, Van de Voorde P, Wang TL, Welsford M, Hazinski MF, Nolan JP, Morley PT; ILCOR Collaborators. 2018 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations Summary. *Resuscitation*. 2018;133:194–206. doi: 10.1016/j.resuscitation.2018.10.017
- Soar J, Donnino MW, Maconochie I, Aickin R, Atkins DL, Andersen LW, Berg KM, Bingham R, Böttiger BW, Callaway CW, Couper K, Couto TB, de Caen AR, Deakin CD, Drennan IR, Guerguerian AM, Lavonas EJ, Meaney PA, Nadkarni VM, Neumar RW, Ng KC, Nicholson TC, Nuthall GA, Ohshimo S, O'Neil BJ, Ong GY, Paiva EF, Parr MJ, Reis AG, Reynolds JC, Ristagno G, Sandroni C, Schexnayder SM, Scholefield BR, Shimizu N, Tijssen JA, Van de Voorde P, Wang TL, Welsford M, Hazinski MF, Nolan JP, Morley PT; ILCOR Collaborators. 2018 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations Summary. *Resuscitation*. 2018;133:e714–e730. doi: 10.1161/CIR.0000000000000611
- Morrison LJ, Deakin CD, Morley PT, Callaway CW, Kerber RE, Kronick SL, Lavonas EJ, Link MS, Neumar RW, Otto CW, et al; on behalf of the Advanced Life Support Chapter Collaborators. Part 8: advanced life support: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2010;122:S345–421. doi: 10.1161/CIRCULATIONAHA.110.971051
- Callaway CW, Soar J, Aibiki M, Böttiger BW, Brooks SC, Deakin CD, Donnino MW, Drajer S, Kloeck W, Morley PT, et al; on behalf of the Advanced Life Support Chapter Collaborators. Part 4: advanced life support: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2015;132(suppl 1):S84–S145. doi: 10.1161/cir.0000000000000273
- Deakin CD, Morrison LJ, Morley PT, Callaway CW, Kerber RE, Kronick SL, Lavonas EJ, Link MS, Neumar RW, Otto CW, et al; on behalf of the Advanced Life Support Chapter Collaborators. Part 8: advanced life support: 2010 International Consensus on Cardiopulmonary

- Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2010;81(suppl 1):e93–e174. doi: 10.1016/j.resuscitation.2010.08.027
9. PRISMA. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) website. <http://www.prisma-statement.org/>. Accessed August 13, 2020.
 10. Schünemann H, Brozek J, Guyatt G, Oxman A, eds. *GRADE Handbook*; 2013. <https://gdt.gradepro.org/app/handbook/handbook.html>. August 13, 2020.
 11. Morley PT, Atkins DL, Finn JC, Maconochie I, Nolan JP, Rabi Y, Singletary EM, Wang TL, Welsford M, Olasveengen TM, et al. Evidence evaluation process and management of potential conflicts of interest: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2020;142(suppl 1):e28–e40. doi: 10.1161/CIR.0000000000000891
 - 11a. Morley PT, Atkins DL, Finn JC, Maconochie I, Nolan JP, Rabi Y, Singletary EM, Wang TL, Welsford M, Olasveengen TM, et al. Evidence evaluation process and management of potential conflicts of interest: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. In press.
 12. Morley PT, Zaritsky A. The evidence evaluation process for the 2005 International Consensus Conference on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation*. 2005;67:167–170. doi: 10.1016/j.resuscitation.2005.09.007
 13. Morley PT, Atkins DL, Billi JE, Bossaert L, Callaway CW, de Caen AR, Deakin CD, Eigel B, Hazinski MF, Hickey RW, et al. Part 3: evidence evaluation process: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2010;122(suppl 2):S283–S290. doi: 10.1161/CIRCULATIONAHA.110.970947
 14. International Liaison Committee on Resuscitation website. <https://www.ilcor.org>. Accessed August 13, 2020.
 15. Perkins GD, Morley PT, Nolan JP, Soar J, Berg K, Olasveengen T, Wyckoff M, Greif R, Singletary N, Castren M, de Caen A, Wang T, Escalante R, Merchant RM, Hazinski M, Kloeck D, Heriot G, Couper K, Neumar R. International Liaison Committee on Resuscitation: COVID-19 consensus on science, treatment recommendations and task force insights. *Resuscitation*. 2020;151:145–147. doi: 10.1016/j.resuscitation.2020.04.035
 16. Neumar RW, Otto CW, Link MS, Kronick SL, Shuster M, Callaway CW, Kudenchuk PJ, Ornato JP, McNally B, Silvers SM, et al. Part 8: adult advanced cardiovascular life support: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2010;122:S729–S767. doi: 10.1161/CIRCULATIONAHA.110.970988
 17. Soar J, Nolan JP, Böttiger BW, Perkins GD, Lott C, Carli P, Pellis T, Sandroni C, Skrifvars MB, Smith GB, Sunde K, Deakin CD; Adult advanced life support section Collaborators. European Resuscitation Council Guidelines for Resuscitation 2015: Section 3. Adult advanced life support. *Resuscitation*. 2015;95:100–147. doi: 10.1016/j.resuscitation.2015.07.016
 18. Otto Q, Deakin C, Morley P, Soar J. Anticipatory manual defibrillator charging during advanced life support: a scoping review. *Resuscitation Plus*. 2020;1–2:100004. <https://doi.org/10.1016/j.resplu.2020.100004>
 19. Mahood Q, Van Eerd D, Irvin E. Searching for grey literature for systematic reviews: challenges and benefits. *Res Synth Methods*. 2014;5:221–234. doi: 10.1002/jrsm.1106
 20. Paez A. Gray literature: An important resource in systematic reviews. *J Evid Based Med*. 2017;10:233–240. doi: 10.1111/jebm.12266
 21. Edelson DP, Robertson-Dick BJ, Yuen TC, Eilevstjønn J, Walsh D, Bareis CJ, Vanden Hoek TL, Abella BS. Safety and efficacy of defibrillator charging during ongoing chest compressions: a multi-center study. *Resuscitation*. 2010;81:1521–1526. doi: 10.1016/j.resuscitation.2010.07.014
 22. Koch Hansen L, Mohammed A, Pedersen M, Folkestad L, Brodersen J, Hey T, Lyhne Christensen N, Carter-Storch R, Bendix K, Hansen MR, Brabrand M. The Stop-Only-While-Shocking algorithm reduces hands-off time by 17% during cardiopulmonary resuscitation - a simulation study. *Eur J Emerg Med*. 2016;23:413–417. doi: 10.1097/MEJ.0000000000000282
 23. Hansen LK, Folkestad L, Brabrand M. Defibrillator charging before rhythm analysis significantly reduces hands-off time during resuscitation: a simulation study. *Am J Emerg Med*. 2013;31:395–400. doi: 10.1016/j.ajem.2012.08.029
 24. Kemper M, Zech A, Lazarovici M, Zwissler B, Prückner S, Meyer O. Defibrillator charging before rhythm analysis causes peri-shock pauses exceeding guideline recommended maximum 5 s: A randomized simulation trial. *Anaesthesist*. 2019;68:546–554. doi: 10.1007/s00101-019-0623-x
 25. Koster RW, Walker RG, Chapman FW. Recurrent ventricular fibrillation during advanced life support care of patients with prehospital cardiac arrest. *Resuscitation*. 2008;78:252–257. doi: 10.1016/j.resuscitation.2008.03.231
 26. Sakai T, Iwami T, Tasaki O, Kawamura T, Hayashi Y, Rinka H, Ohishi Y, Mohri T, Kishimoto M, Nishiuchi T, Kajino K, Matsumoto H, Uejima T, Nitta M, Shiokawa C, Ikeuchi H, Hiraide A, Sugimoto H, Kuwagata Y. Incidence and outcomes of out-of-hospital cardiac arrest with shock-resistant ventricular fibrillation: Data from a large population-based cohort. *Resuscitation*. 2010;81:956–961. doi: 10.1016/j.resuscitation.2010.04.015
 27. Holmén J, Hollenberg J, Claesson A, Herrera MJ, Azeli Y, Herlitz J, Axelsson C. Survival in ventricular fibrillation with emphasis on the number of defibrillations in relation to other factors at resuscitation. *Resuscitation*. 2017;113:33–38. doi: 10.1016/j.resuscitation.2017.01.006
 28. Hasegawa M, Abe T, Nagata T, Onozuka D, Hagihara A. The number of prehospital defibrillation shocks and 1-month survival in patients with out-of-hospital cardiac arrest. *Scand J Trauma Resusc Emerg Med*. 2015;23:34. doi: 10.1186/s13049-015-0112-4
 29. Mapp JG, Hans AJ, Darrington AM, Ross EM, Ho CC, Miramontes DA, Harper SA, Wampler DA; Prehospital Research and Innovation in Military and Expeditionary Environments (PRIME) Research Group. Prehospital double sequential defibrillation: a matched case-control study. *Acad Emerg Med*. 2019;26:994–1001. doi: 10.1111/acem.13672
 30. Ross EM, Redman TT, Harper SA, Mapp JG, Wampler DA, Miramontes DA. Dual defibrillation in out-of-hospital cardiac arrest: A retrospective cohort analysis. *Resuscitation*. 2016;106:14–17. doi: 10.1016/j.resuscitation.2016.06.011
 31. Cortez E, Krebs W, Davis J, Keseg DP, Panchal AR. Use of double sequential external defibrillation for refractory ventricular fibrillation during out-of-hospital cardiac arrest. *Resuscitation*. 2016;108:82–86. doi: 10.1016/j.resuscitation.2016.08.002
 32. Beck LR, Ostermayer DG, Ponce JN, Srinivasan S, Wang HE. Effectiveness of prehospital dual sequential defibrillation for refractory ventricular fibrillation and ventricular tachycardia cardiac arrest. *Prehosp Emerg Care*. 2019;23:597–602. doi: 10.1080/10903127.2019.1584256
 33. Emmerson AC, Whitbread M, Fothergill RT. Double sequential defibrillation therapy for out-of-hospital cardiac arrests: The London experience. *Resuscitation*. 2017;117:97–101. doi: 10.1016/j.resuscitation.2017.06.011
 34. Cabañas JG, Myers JB, Williams JG, De Maio VJ, Bachman MW. Double sequential external defibrillation in out-of-hospital refractory ventricular fibrillation: a report of ten cases. *Prehosp Emerg Care*. 2015;19:126–130. doi: 10.3109/10903127.2014.942476
 35. Cheskes S, Wudwud A, Turner L, McLeod S, Summers J, Morrison LJ, Verbeek PR. The impact of double sequential external defibrillation on termination of refractory ventricular fibrillation during out-of-hospital cardiac arrest. *Resuscitation*. 2019;139:275–281. doi: 10.1016/j.resuscitation.2019.04.038
 36. Deakin CD, Morley P, Soar J, Drennan IR; on behalf of the International Liaison Committee on Resuscitation Advanced Life Support Task Force. Double sequence defibrillation (ALS): systematic review. 2020. <https://costr.ilcor.org/document/header-double-sequence-defibrillation-task-force-systematic-review-costr>. Accessed August 13, 2020.
 37. Cheskes S, Dorian P, Feldman M, McLeod S, Scales DC, Pinto R, Turner L, Morrison LJ, Drennan IR, Verbeek PR. Double sequential external defibrillation for refractory ventricular fibrillation: the DOSE VF pilot randomized controlled trial. *Resuscitation*. 2020;150:178–184. doi: 10.1016/j.resuscitation.2020.02.010
 38. Zijlstra JA, Bekkers LE, Hulleman M, Beesems SG, Koster RW. Automated external defibrillator and operator performance in out-of-hospital cardiac arrest. *Resuscitation*. 2017;118:140–146. doi: 10.1016/j.resuscitation.2017.05.017
 39. Israelsson J, Wangenheim BV, Årestedt K, Semark B, Schildmeijer K, Carlsson J. Sensitivity and specificity of two different automated external defibrillators. *Resuscitation*. 2017;120:108–112. doi: 10.1016/j.resuscitation.2017.09.009
 40. Nehme Z, Andrew E, Nair R, Bernard S, Smith K. Manual versus semiautomatic rhythm analysis and defibrillation for out-of-hospital cardiac arrest. *Circ Cardiovasc Qual Outcomes*. 2017;10 doi: 10.1161/CIRCOUTCOMES.116.003577
 41. Loma-Osorio P, Nunez M, Aboal J, Bosch D, Batlle P, Ruiz de Morales E, Ramos R, Brugada J, Onaga H, Morales A, et al. The Girona Territori Cardioprotegít Project: performance evaluation of public defibrillators. *Rev Esp Cardiol (Engl Ed)*. 2018;71:79–85. doi: 10.1016/j.rec.2017.04.011
 42. Cheskes S, Hillier M, Byers A, Verbeek PR, Drennan IR, Zhan C, Morrison LJ. The association between manual mode defibrillation, pre-shock pause duration and appropriate shock delivery when employed by basic life support paramedics during out-of-hospital cardiac arrest. *Resuscitation*. 2015;90:61–66. doi: 10.1016/j.resuscitation.2015.02.022

43. Sunde K, Jacobs I, Deakin CD, Hazinski MF, Kerber RE, Koster RW, Morrison LJ, Nolan JP, Sayre MR; on behalf of the Defibrillation Chapter Collaborators. Part 6: defibrillation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2010;81(suppl 1):e71–85. doi: 10.1016/j.resuscitation.2010.08.025
44. Jacobs I, Sunde K, Deakin CD, Hazinski MF, Kerber RE, Koster RW, Morrison LJ, Nolan JP, Sayre MR, Defibrillation Chapter C. Part 6: defibrillation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2010;122 (Suppl 2):S325–337. doi: 10.1161/CIRCULATIONAHA.110.971010
45. Freese JP, Jorgenson DB, Liu PY, Innes J, Matallana L, Nammi K, Donohoe RT, Whitbread M, Silverman RA, Prezant DJ. Waveform analysis-guided treatment versus a standard shock-first protocol for the treatment of out-of-hospital cardiac arrest presenting in ventricular fibrillation: results of an international randomized, controlled trial. *Circulation*. 2013;128:995–1002. doi: 10.1161/CIRCULATIONAHA.113.003273
46. Shandilya S, Ward K, Kurz M, Najarian K. Non-linear dynamical signal characterization for prediction of defibrillation success through machine learning. *BMC Med Inform Decis Mak*. 2012;12:116. doi: 10.1186/1472-6947-12-116
47. Nakagawa Y, Sato Y, Kojima T, Wakabayashi T, Morita S, Amino M, Inokuchi S. Electrical defibrillation outcome prediction by waveform analysis of ventricular fibrillation in cardiac arrest out of hospital patients. *Tokai J Exp Clin Med*. 2012;37:1–5.
48. Lin LY, Lo MT, Ko PC, Lin C, Chiang WC, Liu YB, Hu K, Lin JL, Chen WJ, Ma MH. Detrended fluctuation analysis predicts successful defibrillation for out-of-hospital ventricular fibrillation cardiac arrest. *Resuscitation*. 2010;81:297–301. doi: 10.1016/j.resuscitation.2009.12.003
49. Wu X, Bisera J, Tang W. Signal integral for optimizing the timing of defibrillation. *Resuscitation*. 2013;84:1704–1707. doi: 10.1016/j.resuscitation.2013.08.005
50. Coult J, Blackwood J, Sherman L, Rea TD, Kudenchuk PJ, Kwok H. Ventricular fibrillation waveform analysis during chest compressions to predict survival from cardiac arrest. *Circ Arrhythm Electrophysiol*. 2019;12:e006924. doi: 10.1161/CIRCEP.118.006924
51. He M, Lu Y, Zhang L, Zhang H, Gong Y, Li Y. Combining amplitude spectrum area with previous shock information using neural networks improves prediction performance of defibrillation outcome for subsequent shocks in out-of-hospital cardiac arrest patients. *PLoS One*. 2016;11:e0149115. doi: 10.1371/journal.pone.0149115
52. Balderston JR, Gertz ZM, Ellenbogen KA, Schaaf KP, Ornato JP. Association between ventricular fibrillation amplitude immediately prior to defibrillation and defibrillation success in out-of-hospital cardiac arrest. *Am Heart J*. 2018;201:72–76. doi: 10.1016/j.ahj.2018.04.002
53. Agerskov M, Hansen MB, Nielsen AM, Møller TP, Wissenberg M, Rasmussen LS. Return of spontaneous circulation and long-term survival according to feedback provided by automated external defibrillators. *Acta Anaesthesiol Scand*. 2017;61:1345–1353. doi: 10.1111/aa.12992
54. Coult J, Kwok H, Sherman L, Blackwood J, Kudenchuk PJ, Rea TD. Ventricular fibrillation waveform measures combined with prior shock outcome predict defibrillation success during cardiopulmonary resuscitation. *J Electrocardiol*. 2018;51:99–106. doi: 10.1016/j.jelectrocard.2017.07.016
55. Hulleman M, Salcido DD, Menegazzi JJ, Sovereign PC, Tan HL, Blom MT, Koster RW. Predictive value of amplitude spectrum area of ventricular fibrillation waveform in patients with acute or previous myocardial infarction in out-of-hospital cardiac arrest. *Resuscitation*. 2017;120:125–131. doi: 10.1016/j.resuscitation.2017.08.219
56. Jin D, Dai C, Gong Y, Lu Y, Zhang L, Quan W, Li Y. Does the choice of definition for defibrillation and CPR success impact the predictability of ventricular fibrillation waveform analysis? *Resuscitation*. 2017;111:48–54. doi: 10.1016/j.resuscitation.2016.11.022
57. Coult J, Sherman L, Kwok H, Blackwood J, Kudenchuk PJ, Rea TD. Short ECG segments predict defibrillation outcome using quantitative waveform measures. *Resuscitation*. 2016;109:16–20. doi: 10.1016/j.resuscitation.2016.09.020
58. Indik JH, Conover Z, McGovern M, Silver AE, Spaitte DW, Bobrow BJ, Kern KB. Association of amplitude spectral area of the ventricular fibrillation waveform with survival of out-of-hospital ventricular fibrillation cardiac arrest. *J Am Coll Cardiol*. 2014;64:1362–1369. doi: 10.1016/j.jacc.2014.06.1196
59. Hall M, Phelps R, Fahrenbruch C, Sherman L, Blackwood J, Rea TD. Myocardial substrate in secondary ventricular fibrillation: insights from quantitative waveform measures. *Prehosp Emerg Care*. 2011;15:388–392. doi: 10.3109/10903127.2011.561407
60. Foomany FH, Umapathy K, Sugavaneswaran L, Krishnan S, Masse S, Farid T, Nair K, Dorian P, Nanthakumar K. Wavelet-based markers of ventricular fibrillation in optimizing human cardiac resuscitation. *Conf Proc IEEE Eng Med Biol Soc*. 2010;2010:2001–2004. doi: 10.1109/IEMBS.2010.5627841
61. Endoh H, Hida S, Oohashi S, Hayashi Y, Kinoshita H, Honda T. Prompt prediction of successful defibrillation from 1-s ventricular fibrillation waveform in patients with out-of-hospital sudden cardiac arrest. *J Anesth*. 2011;25:34–41. doi: 10.1007/s00540-010-1043-x
62. Aiello S, Perez M, Cogan C, Baetiong A, Miller SA, Radhakrishnan J, Kaufman CL, Gazmuri RJ. Real-time ventricular fibrillation amplitude-spectral area analysis to guide timing of shock delivery improves defibrillation efficacy during cardiopulmonary resuscitation in swine. *J Am Heart Assoc*. 2017;6 doi: 10.1161/JAHA.117.006749
63. Hidano D, Coult J, Blackwood J, Fahrenbruch C, Kwok H, Kudenchuk P, Rea T. Ventricular fibrillation waveform measures and the etiology of cardiac arrest. *Resuscitation*. 2016;109:71–75. doi: 10.1016/j.resuscitation.2016.10.007
64. Howe A, Escalona OJ, Di Maio R, Massot B, Cromie NA, Darragh KM, Adgey J, McEneaney DJ. A support vector machine for predicting defibrillation outcomes from waveform metrics. *Resuscitation*. 2014;85:343–349. doi: 10.1016/j.resuscitation.2013.11.021
65. Nakagawa Y, Amino M, Inokuchi S, Hayashi S, Wakabayashi T, Noda T. Novel CPR system that predicts return of spontaneous circulation from amplitude spectral area before electric shock in ventricular fibrillation. *Resuscitation*. 2017;113:8–12. doi: 10.1016/j.resuscitation.2016.12.025
66. Granfeldt A, Avis SR, Nicholson TC, Holmberg MJ, Moskowitz A, Coker A, Berg KM, Parr MJ, Donnino MW, Soar J, Nation K, Andersen LW; International Liaison Committee on Resuscitation Advanced Life Support Task Force Collaborators. Advanced airway management during adult cardiac arrest: A systematic review. *Resuscitation*. 2019;139:133–143. doi: 10.1016/j.resuscitation.2019.04.003
67. Spindelboeck W, Gemes G, Strasser C, Toescher K, Kores B, Metnitz P, Haas J, Prause G. Arterial blood gases during and their dynamic changes after cardiopulmonary resuscitation: A prospective clinical study. *Resuscitation*. 2016;106:24–29. doi: 10.1016/j.resuscitation.2016.06.013
68. Patel JK, Schoenfeld E, Parikh PB, Parnia S. Association of Arterial Oxygen Tension During In-Hospital Cardiac Arrest With Return of Spontaneous Circulation and Survival. *J Intensive Care Med*. 2018;33:407–414. doi: 10.1177/0885066616658420
69. Rognås L, Hansen TM, Kirkegaard H, Tønnesen E. Standard operating procedure changed pre-hospital critical care anaesthesiologists' behaviour: a quality control study. *Scand J Trauma Resusc Emerg Med*. 2013;21:84. doi: 10.1186/1757-7241-21-84
70. Allen SG, Brewer L, Gillis ES, Pace NL, Sakata DJ, Orr JA. A Turbine-Driven Ventilator Improves Adherence to Advanced Cardiac Life Support Guidelines During a Cardiopulmonary Resuscitation Simulation. *Respir Care*. 2017;62:1166–1170. doi: 10.4187/respcare.05368
71. El Sayed M, Tamim H, Mailhac A, N Clay M. Ventilator use by emergency medical services during 911 calls in the United States. *Am J Emerg Med*. 2018;36:763–768. doi: 10.1016/j.ajem.2017.10.008
72. El Sayed MJ, Tamim H, Mailhac A, Mann NC. Impact of prehospital mechanical ventilation: A retrospective matched cohort study of 911 calls in the United States. *Medicine (Baltimore)*. 2019;98:e13990. doi: 10.1097/MD.00000000000013990
73. Holmberg MJ, Geri G, Wiberg S, Guerguerian AM, Donnino MW, Nolan JP, Deakin CD, Andersen LW; International Liaison Committee on Resuscitation's (ILCOR) Advanced Life Support and Pediatric Task Forces. Extracorporeal cardiopulmonary resuscitation for cardiac arrest: A systematic review. *Resuscitation*. 2018;131:91–100. doi: 10.1016/j.resuscitation.2018.07.029
74. Sutton RM, French B, Meaney PA, Topjian AA, Parshuram CS, Edelson DP, Schexnayder S, Abella BS, Merchant RM, Bembea M, Berg RA, Nadkarni VM; American Heart Association's Get With The Guidelines–Resuscitation Investigators. Physiologic monitoring of CPR quality during adult cardiac arrest: A propensity-matched cohort study. *Resuscitation*. 2016;106:76–82. doi: 10.1016/j.resuscitation.2016.06.018
75. Schnaubelt S, Sulzgruber P, Menger J, Skhirtladze-Dworschak K, Sterz F, Dworschak M. Regional cerebral oxygen saturation during cardiopulmonary resuscitation as a predictor of return of spontaneous circulation and favourable neurological outcome - A review of the current literature. *Resuscitation*. 2018;125:39–47. doi: 10.1016/j.resuscitation.2018.01.028
76. Cournoyer A, Iseppon M, Chauny JM, Denault A, Cossette S, Notebaert É. Near-infrared Spectroscopy Monitoring During Cardiac Arrest: A Systematic Review and Meta-analysis. *Acad Emerg Med*. 2016;23:851–862. doi: 10.1111/acem.12980
77. Prosen G, Strnad M, Doniger SJ, Markota A, Stožer A, Borovnik-Lesjak V, Mekš D. Cerebral tissue oximetry levels during prehospital management

- of cardiac arrest - A prospective observational study. *Resuscitation*. 2018;129:141–145. doi: 10.1016/j.resuscitation.2018.05.014
78. Tsukuda J, Fujitani S, Morisawa K, Shimozawa N, Lohman BD, Okamoto K, Kawaguchi T, Takamatsu Y, Fujii S, Ozaki M, Rahman M, Taira Y. Near-infrared spectroscopy monitoring during out-of-hospital cardiac arrest: can the initial cerebral tissue oxygenation index predict ROSC? *Emerg Med J*. 2019;36:33–38. doi: 10.1136/emered-2018-207533
 79. Yazar MA, Açıkgoz MB, Bayram A. Does chest compression during cardiopulmonary resuscitation provide sufficient cerebral oxygenation? *Turk J Med Sci*. 2019;49:311–317. doi: 10.3906/sag-1809-165
 80. Singer AJ, Nguyen RT, Ravishanker ST, Schoenfeld ER, Thode HC Jr, Henry MC, Parnia S. Cerebral oximetry versus end tidal CO₂ in predicting ROSC after cardiac arrest. *Am J Emerg Med*. 2018;36:403–407. doi: 10.1016/j.ajem.2017.08.046
 81. Nagdyman N, Ewert P, Peters B, Miera O, Fleck T, Berger F. Comparison of different near-infrared spectroscopic cerebral oxygenation indices with central venous and jugular venous oxygenation saturation in children. *Paediatr Anaesth*. 2008;18:160–166. doi: 10.1111/j.1460-9592.2007.02365.x
 82. Kudenchuk PJ, Brown SP, Daya M, Rea T, Nichol G, Morrison LJ, Leroux B, Vaillancourt C, Wittwer L, Callaway CW, Christenson J, Egan D, Ornato JP, Weisfeldt ML, Stiell IG, Idris AH, Aufderheide TP, Dunford JV, Colella MR, Vilke GM, Brienza AM, Desvigne-Nickens P, Gray PC, Gray R, Seals N, Straight R, Dorian P; Resuscitation Outcomes Consortium Investigators. Amiodarone, lidocaine, or placebo in out-of-hospital cardiac arrest. *N Engl J Med*. 2016;374:1711–1722. doi: 10.1056/NEJMoa1514204
 83. Perkins GD, Ji C, Deakin CD, Quinn T, Nolan JP, Scomparin C, Regan S, Long J, Slowther A, Pocock H, Black JJM, Moore F, Fothergill RT, Rees N, O'Shea L, Docherty M, Gunson I, Han K, Charlton K, Finn J, Petrou S, Stallard N, Gates S, Lall R; PARAMEDIC2 Collaborators. A randomized trial of epinephrine in out-of-hospital cardiac arrest. *N Engl J Med*. 2018;379:711–721. doi: 10.1056/NEJMoa1806842
 84. Daya MR, Leroux BG, Dorian P, Rea TD, Newgard CD, Morrison LJ, Lupton JR, Menegazzi JJ, Ornato JP, Sopko G, Christenson J, Idris A, Mody P, Vilke GM, Herdeman C, Barbic D, Kudenchuk PJ; Resuscitation Outcomes Consortium Investigators. Survival after intravenous versus intraosseous amiodarone, lidocaine, or placebo in out-of-hospital shock-refractory cardiac arrest. *Circulation*. 2020;141:188–198. doi: 10.1161/CIRCULATIONAHA.119.042240
 85. Nolan JP, Deakin CD, Ji C, Gates S, Rosser A, Lall R, Perkins GD. Intraosseous versus intravenous administration of adrenaline in patients with out-of-hospital cardiac arrest: a secondary analysis of the PARAMEDIC2 placebo-controlled trial. *Intensive Care Med*. 2020;46:954–962. doi: 10.1007/s00134-019-05920-7
 86. Holmberg MJ, Issa MS, Moskowitz A, Morley P, Welsford M, Neumar RW, Paiva EF, Coker A, Hansen CK, Andersen LW, Donnino MW, Berg KM; International Liaison Committee on Resuscitation Advanced Life Support Task Force Collaborators. Vasopressors during adult cardiac arrest: A systematic review and meta-analysis. *Resuscitation*. 2019;139:106–121. doi: 10.1016/j.resuscitation.2019.04.008
 87. Ali MU, Fitzpatrick-Lewis D, Kenny M, Raina P, Atkins DL, Soar J, Nolan J, Ristagno G, Sherifali D. Effectiveness of antiarrhythmic drugs for shockable cardiac arrest: A systematic review. *Resuscitation*. 2018;132:63–72. doi: 10.1016/j.resuscitation.2018.08.025
 88. Granfeldt A, Avis SR, Lind PC, Holmberg MJ, Kleinman M, Maconochiel, Hsu CH, Fernanda de Almeida M, Wang TL, Neumar RW, Andersen LW. Intravenous vs. intraosseous administration of drugs during cardiac arrest: A systematic review. *Resuscitation*. 2020;149:150–157. doi: 10.1016/j.resuscitation.2020.02.025
 89. Mody P, Brown SP, Kudenchuk PJ, Chan PS, Khera R, Ayers C, Pandey A, Kern KB, de Lemos JA, Link MS, Idris AH. Intraosseous versus intravenous access in patients with out-of-hospital cardiac arrest: Insights from the resuscitation outcomes consortium continuous chest compression trial. *Resuscitation*. 2019;134:69–75. doi: 10.1016/j.resuscitation.2018.10.031
 90. Feinstein BA, Stubbs BA, Rea T, Kudenchuk PJ. Intraosseous compared to intravenous drug resuscitation in out-of-hospital cardiac arrest. *Resuscitation*. 2017;117:91–96. doi: 10.1016/j.resuscitation.2017.06.014
 91. Kawano T, Grunau B, Scheuermeyer FX, Gibo K, Fordyce CB, Lin S, Stenstrom R, Schlamp R, Jenneson S, Christenson J. Intraosseous vascular access is associated with lower survival and neurologic recovery among patients with out-of-hospital cardiac arrest. *Ann Emerg Med*. 2018;71:588–596. doi: 10.1016/j.annemergmed.2017.11.015
 92. Zhang Y, Zhu J, Liu Z, Gu L, Zhang W, Zhan H, Hu C, Liao J, Xiong Y, Idris AH. Intravenous versus intraosseous adrenaline administration in out-of-hospital cardiac arrest: a retrospective cohort study. *Resuscitation*. 2020;149:209–216. doi: 10.1016/j.resuscitation.2020.01.009
 93. Link MS, Berkow LC, Kudenchuk PJ, Halperin HR, Hess EP, Moitra VK, Neumar RW, O'Neil BJ, Paxton JH, Silvers SM, et al. Part 7: adult advanced cardiovascular life support: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2015;132(suppl 2):S444–S464. doi: 10.1161/CIR.0000000000000261
 94. Tsai MS, Chuang PY, Huang CH, Tang CH, Yu PH, Chang WT, Chen WJ. Postarrest steroid use may improve outcomes of cardiac arrest survivors. *Crit Care Med*. 2019;47:167–175. doi: 10.1097/CCM.0000000000003468
 95. Niimura T, Zamami Y, Koyama T, Izawa-Ishizawa Y, Miyake M, Koga T, Harada K, Ohshima A, Imai T, Kondo Y, Imanishi M, Takechi K, Fukushima K, Horinouchi Y, Ikeda Y, Fujino H, Tsuchiya K, Tamaki T, Hinotsu S, Kano MR, Ishizawa K. Hydrocortisone administration was associated with improved survival in Japanese patients with cardiac arrest. *Sci Rep*. 2017;7:17919. doi: 10.1038/s41598-017-17686-3
 96. Ahn S, Kim YJ, Sohn CH, Seo DW, Lim KS, Donnino MW, Kim WY. Sodium bicarbonate on severe metabolic acidosis during prolonged cardiopulmonary resuscitation: a double-blind, randomized, placebo-controlled pilot study. *J Thorac Dis*. 2018;10:2295–2302. doi: 10.21037/jtd.2018.03.124
 97. Chen YC, Hung MS, Liu CY, Hsiao CT, Yang YH. The association of emergency department administration of sodium bicarbonate after out of hospital cardiac arrest with outcomes. *Am J Emerg Med*. 2018;36:1998–2004. doi: 10.1016/j.ajem.2018.03.010
 98. Wang CH, Huang CH, Chang WT, Tsai MS, Yu PH, Wu YW, Hung KY, Chen WJ. The effects of calcium and sodium bicarbonate on severe hyperkalaemia during cardiopulmonary resuscitation: A retrospective cohort study of adult in-hospital cardiac arrest. *Resuscitation*. 2016;98:105–111. doi: 10.1016/j.resuscitation.2015.09.384
 99. Kawano T, Grunau B, Scheuermeyer FX, Gibo K, Dick W, Fordyce CB, Dorian P, Stenstrom R, Straight R, Christenson J. Prehospital sodium bicarbonate use could worsen long term survival with favorable neurological recovery among patients with out-of-hospital cardiac arrest. *Resuscitation*. 2017;119:63–69. doi: 10.1016/j.resuscitation.2017.08.008
 100. Kim J, Kim K, Park J, Jo YH, Lee JH, Hwang JE, Ha C, Ko YS, Jung E. Sodium bicarbonate administration during ongoing resuscitation is associated with increased return of spontaneous circulation. *Am J Emerg Med*. 2016;34:225–229. doi: 10.1016/j.ajem.2015.10.037
 101. Reynolds JC, Issa MS, Nicholson T, Drennan IR, Berg K, O'Neil BJ, Welsford M; on behalf of the Advanced Life Support Task Force of the International Liaison Committee on Resuscitation. Prognostication with point-of-care echocardiography during cardiac arrest: a systematic review. *Resuscitation*. 2020;152:56–68. doi: https://doi.org/10.1016/j.resuscitation.2020.05.004
 102. Lien WC, Hsu SH, Chong KM, Sim SS, Wu MC, Chang WT, Fang CC, Ma MH, Chen SC, Chen WJ. US-CAB protocol for ultrasonographic evaluation during cardiopulmonary resuscitation: Validation and potential impact. *Resuscitation*. 2018;127:125–131. doi: 10.1016/j.resuscitation.2018.01.051
 103. Salen P, Melniker L, Chooljian C, Rose JS, Altevveer J, Reed J, Heller M. Does the presence or absence of sonographically identified cardiac activity predict resuscitation outcomes of cardiac arrest patients? *Am J Emerg Med*. 2005;23:459–462. doi: 10.1016/j.ajem.2004.11.007
 104. Salen P, O'Connor R, Sierzynski P, Passarello B, Pancu D, Melanson S, Arcona S, Reed J, Heller M. Can cardiac sonography and capnography be used independently and in combination to predict resuscitation outcomes? *Acad Emerg Med*. 2001;8:610–615. doi: 10.1111/j.1553-2712.2001.tb00172.x
 105. Tayal VS, Kline JA. Emergency echocardiography to detect pericardial effusion in patients in PEA and near-PEA states. *Resuscitation*. 2003;59:315–318. doi: 10.1016/s0300-9572(03)00245-4
 106. Varriale P, Maldonado JM. Echocardiographic observations during in hospital cardiopulmonary resuscitation. *Crit Care Med*. 1997;25:1717–1720. doi: 10.1097/00003246-199710000-00023
 107. Aichinger G, Zechner PM, Prause G, Sacherer F, Wildner G, Anderson CL, Pocalnik M, Wiesspeiner U, Fox JC. Cardiac movement identified on prehospital echocardiography predicts outcome in cardiac arrest patients. *Prehosp Emerg Care*. 2012;16:251–255. doi: 10.3109/10903127.2011.640414
 108. Atkinson PR, Beckett N, French J, Banerjee A, Fraser J, Lewis D. Does point-of-care ultrasound use impact resuscitation length, rates of intervention, and clinical outcomes during cardiac arrest? a study from the Sonography in Hypotension and Cardiac Arrest in the Emergency Department (SHoC-ED) Investigators. *Cureus*. 2019;11:e4456. doi: 10.7759/cureus.4456

109. Blaivas M, Fox JC. Outcome of cardiac arrest patients found to have cardiac standstill on the bedside emergency department echocardiogram. *Acad Emerg Med*. 2001;8:616–621. doi: 10.1111/j.1553-2712.2001.tb00174.x
110. Breitkreutz R, Price S, Steiger HV, Seeger FH, Ilper H, Ackermann H, Rudolph M, Uddin S, Weigand MA, Müller E, Walcher F; Emergency Ultrasound Working Group of the Johann Wolfgang Goethe-University Hospital, Frankfurt am Main. Focused echocardiographic evaluation in life support and peri-resuscitation of emergency patients: a prospective trial. *Resuscitation*. 2010;81:1527–1533. doi: 10.1016/j.resuscitation.2010.07.013
111. Chardoli M, Heidari F, Rabiee H, Sharif-Alhoseini M, Shokoohi H, Rahimi-Movaghar V. Echocardiography integrated ACLS protocol versus conventional cardiopulmonary resuscitation in patients with pulseless electrical activity cardiac arrest. *Chin J Traumatol*. 2012;15:284–287.
112. Chua MT, Chan GW, Kuan WS. Reversible causes in cardiovascular collapse at the emergency department using ultrasonography (REVIVE-US). *Ann Acad Med Singapore*. 2017;46:310–316.
113. Flato UA, Paiva EF, Carballo MT, Buehler AM, Marco R, Timerman A. Echocardiography for prognostication during the resuscitation of intensive care unit patients with non-shockable rhythm cardiac arrest. *Resuscitation*. 2015;92:1–6. doi: 10.1016/j.resuscitation.2015.03.024
114. Gaspari R, Weekes A, Adhikari S, Noble VE, Nomura JT, Theodoro D, Woo M, Atkinson P, Behar D, Brown SM, Caffery T, Douglass E, Fraser J, Haines C, Lam S, Lanspa M, Lewis M, Liebmann O, Limkakeng A, Lopez F, Platz E, Mendoza M, Minnigan H, Moore C, Novik J, Rang L, Scruggs W, Raio C. Emergency department point-of-care ultrasound in out-of-hospital and in-ED cardiac arrest. *Resuscitation*. 2016;109:33–39. doi: 10.1016/j.resuscitation.2016.09.018
115. Kim HB, Suh JY, Choi JH, Cho YS. Can serial focussed echocardiographic evaluation in life support (FEEL) predict resuscitation outcome or termination of resuscitation (TOR)? A pilot study. *Resuscitation*. 2016;101:21–26. doi: 10.1016/j.resuscitation.2016.01.013
116. Zengin S, Yavuz E, Al B, Cindoruk Ş, Altunbaş G, Gümüşboğa H, Yıldırım C. Benefits of cardiac sonography performed by a non-expert sonographer in patients with non-traumatic cardiopulmonary arrest. *Resuscitation*. 2016;102:105–109. doi: 10.1016/j.resuscitation.2016.02.025
117. Querellou E, Leyral J, Brun C, Lévy D, Bessereau J, Meyran D, Le Dreff P. [In and out-of-hospital cardiac arrest and echography: a review]. *Ann Fr Anesth Reanim*. 2009;28:769–778. doi: 10.1016/j.annfar.2009.06.020
118. Blanco P, Volpicelli G. Common pitfalls in point-of-care ultrasound: a practical guide for emergency and critical care physicians. *Crit Ultrasound J*. 2016;8:15. doi: 10.1186/s13089-016-0052-x
119. Aagaard R, Granfeldt A, Bøtker MT, Mygind-Klausen T, Kirkegaard H, Løfgren B. The right ventricle is dilated during resuscitation from cardiac arrest caused by hypovolemia: a porcine ultrasound study. *Crit Care Med*. 2017;45:e963–e970. doi: 10.1097/CCM.0000000000002464
120. Clattenburg EJ, Wroe P, Brown S, Gardner K, Losonczy L, Singh A, Nagdev A. Point-of-care ultrasound use in patients with cardiac arrest is associated prolonged cardiopulmonary resuscitation pauses: A prospective cohort study. *Resuscitation*. 2018;122:65–68. doi: 10.1016/j.resuscitation.2017.11.056
121. Huis In 't Veld MA, Allison MG, Bostick DS, Fisher KR, Goloubeva OG, Witting MD, Winters ME. Ultrasound use during cardiopulmonary resuscitation is associated with delays in chest compressions. *Resuscitation*. 2017;119:95–98. doi: 10.1016/j.resuscitation.2017.07.021
122. Clattenburg EJ, Wroe PC, Gardner K, Schultz C, Gelber J, Singh A, Nagdev A. Implementation of the Cardiac Arrest Sonographic Assessment (CASA) protocol for patients with cardiac arrest is associated with shorter CPR pulse checks. *Resuscitation*. 2018;131:69–73. doi: 10.1016/j.resuscitation.2018.07.030
123. Finn TE, Ward JL, Wu CT, Giles A, Manivel V. COACHRED: A protocol for the safe and timely incorporation of focused echocardiography into the rhythm check during cardiopulmonary resuscitation. *Emerg Med Australas*. 2019;31:1115–1118. doi: 10.1111/1742-6723.13374
124. Paiva EF, Paxton JH, O'Neil BJ. The use of end-tidal carbon dioxide (ETCO₂) measurement to guide management of cardiac arrest: A systematic review. *Resuscitation*. 2018;123:1–7. doi: 10.1016/j.resuscitation.2017.12.003
125. Pearce AK, Davis DP, Minokadeh A, Sell RE. Initial end-tidal carbon dioxide as a prognostic indicator for inpatient PEA arrest. *Resuscitation*. 2015;92:77–81. doi: 10.1016/j.resuscitation.2015.04.025
126. Poon KM, Lui CT, Tsui KL. Prognostication of out-of-hospital cardiac arrest patients by 3-min end-tidal capnometry level in emergency department. *Resuscitation*. 2016;102:80–84. doi: 10.1016/j.resuscitation.2016.02.021
127. Poppe M, Stratil P, Clodi C, Schrieffl C, Nürnberg A, Magnet I, Warenits AM, Hubner P, Lobmeyr E, Schober A, Zajicek A, Testori C. Initial end-tidal carbon dioxide as a predictive factor for return of spontaneous circulation in nonshockable out-of-hospital cardiac arrest patients: A retrospective observational study. *Eur J Anaesthesiol*. 2019;36:524–530. doi: 10.1097/EJA.0000000000000999
128. Wang AY, Huang CH, Chang WT, Tsai MS, Wang CH, Chen WJ. Initial end-tidal CO₂ partial pressure predicts outcomes of in-hospital cardiac arrest. *Am J Emerg Med*. 2016;34:2367–2371. doi: 10.1016/j.ajem.2016.08.052
129. Akinci E, Ramadan H, Yuzbasioglu Y, Coskun F. Comparison of end-tidal carbon dioxide levels with cardiopulmonary resuscitation success presented to emergency department with cardiopulmonary arrest. *Pak J Med Sci*. 2014;30:16–21. doi: 10.12669/pjms.301.4024
130. Böttiger BW, Arntz HR, Chamberlain DA, Bluhmki E, Belmans A, Danays T, Carli PA, Adgey JA, Bode C, Wenzel V; TROICA Trial Investigators; European Resuscitation Council Study Group. Thrombolysis during resuscitation for out-of-hospital cardiac arrest. *N Engl J Med*. 2008;359:2651–2662. doi: 10.1056/NEJMoa070570
131. Javaudin F, Lascarrou JB, Le Bastard Q, Bourry Q, Latour C, De Carvalho H, Le Conte P, Escutnaire J, Hubert H, Montassier E, Leclère B; Research Group of the French National Out-of-Hospital Cardiac Arrest Registry (GR-RéAC). Thrombolysis during resuscitation for out-of-hospital cardiac arrest caused by pulmonary embolism increases 30-day survival: findings from the French National Cardiac Arrest Registry. *Chest*. 2019;156:1167–1175. doi: 10.1016/j.chest.2019.07.015
132. Yousuf T, Brinton T, Ahmed K, Iskander J, Woznicka D, Kramer J, Kopic A, Chadaga AR, Ortiz K. Tissue Plasminogen Activator Use in Cardiac Arrest Secondary to Fulminant Pulmonary Embolism. *J Clin Med Res*. 2016;8:190–195. doi: 10.14740/jocmr2452w
133. Kürkçyan I, Meron G, Sterz F, Janata K, Domanovits H, Holzer M, Berzlanovich A, Bankl HC, Laggner AN. Pulmonary embolism as a cause of cardiac arrest: presentation and outcome. *Arch Intern Med*. 2000;160:1529–1535. doi: 10.1001/archinte.160.10.1529
134. Janata K, Holzer M, Kürkçyan I, Losert H, Riedmüller E, Pikula B, Laggner AN, Laczika K. Major bleeding complications in cardiopulmonary resuscitation: the place of thrombolytic therapy in cardiac arrest due to massive pulmonary embolism. *Resuscitation*. 2003;57:49–55. doi: 10.1016/s0300-9572(02)00430-6
135. Konstantinov IE, Saxena P, Koniuszko MD, Alvarez J, Newman MA. Acute massive pulmonary embolism with cardiopulmonary resuscitation: management and results. *Tex Heart Inst J*. 2007;34:41–5; discussion 45.
136. Doerge HC, Schoendube FA, Loeser H, Walter M, Messmer BJ. Pulmonary embolism: review of a 15-year experience and role in the age of thrombolytic therapy. *Eur J Cardiothorac Surg*. 1996;10:952–957. doi: 10.1016/s1010-7940(96)80396-4
137. Fava M, Loyola S, Bertoni H, Dougnac A. Massive pulmonary embolism: percutaneous mechanical thrombectomy during cardiopulmonary resuscitation. *J Vasc Interv Radiol*. 2005;16:119–123. doi: 10.1097/01.RVI.0000146173.85401.BA
138. Beckett VA, Knight M, Sharpe P. The CAPS Study: incidence, management and outcomes of cardiac arrest in pregnancy in the UK: a prospective, descriptive study. *BJOG*. 2017;124:1374–1381. doi: 10.1111/1471-0528.14521
139. Biderman P, Carmi U, Setton E, Fainblut M, Bachar O, Einav S. Maternal salvage with extracorporeal life support: lessons learned in a single center. *Anesth Analg*. 2017;125:1275–1280. doi: 10.1213/ANE.0000000000002262
140. Schaap TP, Overtoom E, van den Akker T, Zwart JJ, van Roosmalen J, Bloemenkamp KWM. Maternal cardiac arrest in the Netherlands: A nationwide surveillance study. *Eur J Obstet Gynecol Reprod Biol*. 2019;237:145–150. doi: 10.1016/j.ejogrb.2019.04.028
141. Maurin O, Lemoine S, Jost D, Lanoë V, Renard A, Travers S, The Paris Fire Brigade Cardiac Arrest Work Group, Lapostolle F, Tourtier JP. Maternal out-of-hospital cardiac arrest: A retrospective observational study. *Resuscitation*. 2019;135:205–211. doi: 10.1016/j.resuscitation.2018.11.001
142. Kobori S, Toshimitsu M, Nagaoka S, Yaegashi N, Murotsuki J. Utility and limitations of perimortem cesarean section: A nationwide survey in Japan. *J Obstet Gynaecol Res*. 2019;45:325–330. doi: 10.1111/jog.13819
- 142a. Greif R, Bhanji F, Bigham BL, Bray J, Breckwoldt J, Cheng A, Duff JP, Gilfoyle E, Hsieh M-J, Iwami T, et al; on behalf of the Education, Implementation, and Teams Collaborators. Education, implementation, and teams: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2020;142(suppl 1):S222–S283. doi: 10.1161/CIR.0000000000000896

- 142b. Greif R, Bhanji F, Bigham BL, Bray J, Breckwoldt J, Cheng A, Duff JP, Gilfoyle E, Hsieh M-J, Iwami T, et al; on behalf of the Education, Implementation, and Teams Collaborators. Education, implementation, and teams: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. In press.
143. Andersen LW, Lind PC, Vammen L, Høybye M, Holmberg MJ, Granfeldt A. Adult post-cardiac arrest interventions: An overview of randomized clinical trials. *Resuscitation*. 2020;147:1–11. doi: 10.1016/j.resuscitation.2019.12.003
144. Holmberg MJ, Nicholson T, Nolan JP, Schexnayder S, Reynolds J, Nation K, Welsford M, Morley P, Soar J, Berg KM. Oxygenation and ventilation targets after cardiac arrest: a systematic review and meta-analysis. *Resuscitation*. 2020;152:107–115 doi: 10.1016/j.resuscitation.2020.04.031
145. Mackle D, Bellomo R, Bailey M, Beasley R, Deane A, Eastwood G, Finfer S, Freebairn R, King V, Linke N, et al; and the ICU-ROX Investigators the Australian New Zealand Intensive Care Society Clinical Trials Group. Conservative oxygen therapy during mechanical ventilation in the ICU. *N Engl J Med*. 2020;382:989–998. doi: 10.1056/NEJMoa1903297
146. Janz DR, Hollenbeck RD, Pollock JS, McPherson JA, Rice TW. Hyperoxia is associated with increased mortality in patients treated with mild therapeutic hypothermia after sudden cardiac arrest. *Crit Care Med*. 2012;40:3135–3139. doi: 10.1097/CCM.0b013e3182656976
147. Elmer J, Scutella M, Pullalarevu R, Wang B, Vaghasia N, Trzeciak S, Rosario-Rivera BL, Guyette FX, Rittenberger JC, Dezfulian C; Pittsburgh Post-Cardiac Arrest Service (PCAS). The association between hyperoxia and patient outcomes after cardiac arrest: analysis of a high-resolution database. *Intensive Care Med*. 2015;41:49–57. doi: 10.1007/s00134-014-3555-6
148. Roberts BW, Kilgannon JH, Hunter BR, Puskari MA, Pierce L, Donnino M, Leary M, Kline JA, Jones AE, Shapiro NI, Abella BS, Trzeciak S. Association between early hyperoxia exposure after resuscitation from cardiac arrest and neurological disability: prospective multicenter protocol-directed cohort study. *Circulation*. 2018;137:2114–2124. doi: 10.1161/CIRCULATIONAHA.117.032054
149. Wang HE, Prince DK, Drennan IR, Grunau B, Carlbom DJ, Johnson N, Hansen M, Elmer J, Christenson J, Kudenchuk P, Aufderheide T, Weisfeldt M, Idris A, Trzeciak S, Kurz M, Rittenberger JC, Griffiths D, Jasti J, May S; Resuscitation Outcomes Consortium (ROC) Investigators. Post-resuscitation arterial oxygen and carbon dioxide and outcomes after out-of-hospital cardiac arrest. *Resuscitation*. 2017;120:113–118. doi: 10.1016/j.resuscitation.2017.08.244
150. Johnson NJ, Dodampahala K, Rosselot B, Perman SM, Mikkelsen ME, Goyal M, Gaieski DF, Grossestreuer AV. The association between arterial oxygen tension and neurological outcome after cardiac arrest. *Ther Hypothermia Temp Manag*. 2017;7:36–41. doi: 10.1089/ther.2016.0015
151. Humaloja J, Litonius E, Efendijev I, Folger D, Raj R, Pekkarinen PT, Skrifvars MB. Early hyperoxemia is not associated with cardiac arrest outcome. *Resuscitation*. 2019;140:185–193. doi: 10.1016/j.resuscitation.2019.04.035
152. von Auenmueller KI, Christ M, Sasko BM, Trappe HJ. The value of arterial blood gas parameters for prediction of mortality in survivors of out-of-hospital cardiac arrest. *J Emerg Trauma Shock*. 2017;10:134–139. doi: 10.4103/JETS.JETS_146_16
153. Ebner F, Ullén S, Åneman A, Cronberg T, Mattsson N, Friberg H, Hassager C, Kjærgaard J, Kuiper M, Pelosi P, Undén J, Wise MP, Wetterslev J, Nielsen N. Associations between partial pressure of oxygen and neurological outcome in out-of-hospital cardiac arrest patients: an explorative analysis of a randomized trial. *Crit Care*. 2019;23:30. doi: 10.1186/s13054-019-2322-z
154. Eastwood GM, Tanaka A, Espinoza ED, Peck L, Young H, Mårtensson J, Zhang L, Glassford NJ, Hsiao YF, Suzuki S, Bellomo R. Conservative oxygen therapy in mechanically ventilated patients following cardiac arrest: A retrospective nested cohort study. *Resuscitation*. 2016;101:108–114. doi: 10.1016/j.resuscitation.2015.11.026
155. Vaahersalo J, Bendel S, Reinikainen M, Kurolo J, Tiainen M, Raj R, Pettilä V, Varpula T, Skrifvars MB; FINNRESUSCI Study Group. Arterial blood gas tensions after resuscitation from out-of-hospital cardiac arrest: associations with long-term neurologic outcome. *Crit Care Med*. 2014;42:1463–1470. doi: 10.1097/CCM.0000000000000228
156. Jakkula P, Reinikainen M, Hästbacka K, Loisa P, Tiainen M, Pettilä V, Toppila J, Lähde M, Bäcklund M, Okkonen M, et al; and the COMA-CARE study group. Targeting two different levels of both arterial carbon dioxide and arterial oxygen after cardiac arrest and resuscitation: a randomised pilot trial. *Intensive Care Med*. 2018;44:2112–2121. doi: 10.1007/s00134-018-5453-9
157. Young P, Bailey M, Bellomo R, Bernard S, Dicker B, Freebairn R, Henderson S, Mackle D, McArthur C, McGuinness S, Smith T, Swain A, Weatherall M, Beasley R. HyperOxIC Therapy OR NormOxIC Therapy after out-of-hospital cardiac arrest (HOT OR NOT): a randomised controlled feasibility trial. *Resuscitation*. 2014;85:1686–1691. doi: 10.1016/j.resuscitation.2014.09.011
158. Kuisma M, Boyd J, Voipio V, Alasää A, Roine RO, Rosenberg P. Comparison of 30 and the 100% inspired oxygen concentrations during early post-resuscitation period: a randomised controlled pilot study. *Resuscitation*. 2006;69:199–206. doi: 10.1016/j.resuscitation.2005.08.010
159. Bray JE, Hein C, Smith K, Stephenson M, Grantham H, Finn J, Stub D, Cameron P, Bernard S; EXACT Investigators. Oxygen titration after resuscitation from out-of-hospital cardiac arrest: A multi-centre, randomised controlled pilot study (the EXACT pilot trial). *Resuscitation*. 2018;128:211–215. doi: 10.1016/j.resuscitation.2018.04.019
160. Thomas M, Voss S, Bengler J, Kirby K, Nolan JP. Cluster randomised comparison of the effectiveness of 100% oxygen versus titrated oxygen in patients with a sustained return of spontaneous circulation following out of hospital cardiac arrest: a feasibility study. PROXY: post ROSC OXYgenation study. *BMC Emerg Med*. 2019;19:16. doi: 10.1186/s12873-018-0214-1
161. Sekhon MS, Griesdale DE. Individualized perfusion targets in hypoxic ischemic brain injury after cardiac arrest. *Crit Care*. 2017;21:259. doi: 10.1186/s13054-017-1832-9
162. Eastwood GM, Schneider AG, Suzuki S, Peck L, Young H, Tanaka A, Mårtensson J, Warrillow S, McGuinness S, Parke R, Gilder E, McCarthy L, Galt P, Taori G, Elliott S, Lamac T, Bailey M, Harley N, Barge D, Hodgson CL, Morganti-Kossmann MC, Pébay A, Conquest A, Archer JS, Bernard S, Stub D, Hart GK, Bellomo R. Targeted therapeutic mild hypercapnia after cardiac arrest: A phase II multi-centre randomised controlled trial (the CCC trial). *Resuscitation*. 2016;104:83–90. doi: 10.1016/j.resuscitation.2016.03.023
163. Hope Kilgannon J, Hunter BR, Puskari MA, Shea L, Fuller BM, Jones C, Donnino M, Kline JA, Jones AE, Shapiro NI, Abella BS, Trzeciak S, Roberts BW. Partial pressure of arterial carbon dioxide after resuscitation from cardiac arrest and neurological outcome: A prospective multi-center protocol-directed cohort study. *Resuscitation*. 2019;135:212–220. doi: 10.1016/j.resuscitation.2018.11.015
164. Roberts BW, Kilgannon JH, Chansky ME, Mittal N, Wooden J, Trzeciak S. Association between postresuscitation partial pressure of arterial carbon dioxide and neurological outcome in patients with post-cardiac arrest syndrome. *Circulation*. 2013;127:2107–2113. doi: 10.1161/CIRCULATIONAHA.112.000168
165. Ebner F, Harmon MBA, Aneman A, Cronberg T, Friberg H, Hassager C, Juffermans N, Kjærgaard J, Kuiper M, Mattsson N, Pelosi P, Ullén S, Undén J, Wise MP, Nielsen N. Carbon dioxide dynamics in relation to neurological outcome in resuscitated out-of-hospital cardiac arrest patients: an exploratory Target Temperature Management Trial substudy. *Crit Care*. 2018;22:196. doi: 10.1186/s13054-018-2119-5
166. Ameloot K, De Deyne C, Eertmans V, Ferdinande B, Dupont M, Palmers PJ, Petit T, Nuyens P, Maeremans J, Vundelinckx J, Vanhaverbeke M, Belmans A, Peeters R, Demaerel P, Lemmens R, Dens J, Janssens S. Early goal-directed haemodynamic optimization of cerebral oxygenation in comatose survivors after cardiac arrest: the Neuroprotect post-cardiac arrest trial. *Eur Heart J*. 2019;40:1804–1814. doi: 10.1093/eurheartj/ehz120
167. Mentzelopoulos SD, Malachias S, Chamos C, Konstantopoulos D, Ntaidou T, Papastylianou A, Kolliantzaki I, Theodoridi M, Ischaki H, Makris D, Zakyntinos E, Zintzaras E, Sourlas S, Aloizos S, Zakyntinos SG. Vasopressin, steroids, and epinephrine and neurologically favorable survival after in-hospital cardiac arrest: a randomized clinical trial. *JAMA*. 2013;310:270–279. doi: 10.1001/jama.2013.7832
168. Mentzelopoulos SD, Zakyntinos SG, Tzoufi M, Katsios N, Papastylianou A, Gkisioti S, Stathopoulos A, Kollintza A, Stamataki E, Roussos C. Vasopressin, epinephrine, and corticosteroids for in-hospital cardiac arrest. *Arch Intern Med*. 2009;169:15–24. doi: 10.1001/archinternmed.2008.509
169. Donnino MW, Andersen LW, Berg KM, Chase M, Sherwin R, Smithline H, Carney E, Ngo L, Patel PV, Liu X, Cutlip D, Zimetbaum P, Cocchi MN; Collaborating Authors from the Beth Israel Deaconess Medical Center's Center for Resuscitation Science Research Group. Corticosteroid therapy in refractory shock following cardiac arrest: a randomized, double-blind, placebo-controlled trial. *Crit Care*. 2016;20:82. doi: 10.1186/s13054-016-1257-x
170. Nielsen N, Wetterslev J, Cronberg T, Erlinge D, Gasche Y, Hassager C, Horn J, Hovdenes J, Kjærgaard J, Kuiper M, Pellis T, Ståmmet P, Wanscher M, Wise MP, Åneman A, Al-Subaie N, Boesgaard S, Bro-Jeppesen J, Brunetti I, Bugge JF, Hingston CD, Juffermans NP, Koopmans M, Køber L, Langørgen J, Lilja G, Møller JE, Rundgren M, Rylander C, Smid O, Werer C, Winkel P,

- Friberg H; TTM Trial Investigators. Targeted temperature management at 33°C versus 36°C after cardiac arrest. *N Engl J Med*. 2013;369:2197–2206. doi: 10.1056/NEJMoa1310519
171. François B, Cariou A, Clere-Jehl R, Dequin PF, Renon-Carron F, Daix T, Guittion C, Deye N, Legriel S, Plantefève G, Quenot JP, Desachy A, Kamel T, Bedon-Cardé S, Diehl JL, Chudeau N, Karam E, Durand-Zaleski I, Giraudeau B, Vignon P, Le Gouge A; CRICS-TRIGGERSEP Network and the ANTHARTIC Study Group. Prevention of early ventilator-associated pneumonia after cardiac arrest. *N Engl J Med*. 2019;381:1831–1842. doi: 10.1056/NEJMoa1812379
 172. Couper K, Laloo R, Field R, Perkins GD, Thomas M, Yeung J. Prophylactic antibiotic use following cardiac arrest: A systematic review and meta-analysis. *Resuscitation*. 2019;141:166–173. doi: 10.1016/j.resuscitation.2019.04.047
 173. Schünemann HJ, Wiercioch W, Brozek J, Etzeandia-Ikobaltzeta I, Mustafa RA, Manja V, Brignardello-Petersen R, Neumann I, Falavigna M, Alhazzani W, Santesso N, Zhang Y, Meerpohl JJ, Morgan RL, Rochwerf B, Darzi A, Rojas MX, Carrasco-Labra A, Adi Y, AlRayees Z, Riva J, Bollig C, Moore A, Yepes-Nuñez JJ, Cuello C, Waziry R, Akl EA. GRADE Evidence to Decision (ETD) frameworks for adoption, adaptation, and de novo development of trustworthy recommendations: GRADE-ADOLPMENT. *J Clin Epidemiol*. 2017;81:101–110. doi: 10.1016/j.jclinepi.2016.09.009
 174. Ribaric SF, Turel M, Knafelj R, Gorjup V, Stanic R, Gradisek P, Cerovic O, Mirkovic T, Noc M. Prophylactic versus clinically-driven antibiotics in comatose survivors of out-of-hospital cardiac arrest—A randomized pilot study. *Resuscitation*. 2017;111:103–109. doi: 10.1016/j.resuscitation.2016.11.025
 175. Tagami T, Matsui H, Kuno M, Moroe Y, Kaneko J, Unemoto K, Fushimi K, Yasunaga H. Early antibiotics administration during targeted temperature management after out-of-hospital cardiac arrest: a nationwide database study. *BMC Anesthesiol*. 2016;16:89. doi: 10.1186/s12871-016-0257-3
 176. Davies KJ, Walters JH, Kerslake IM, Greenwood R, Thomas MJ. Early antibiotics improve survival following out-of hospital cardiac arrest. *Resuscitation*. 2013;84:616–619. doi: 10.1016/j.resuscitation.2012.11.004
 177. Perbet S, Mongardon N, Dumas F, Bruel C, Lemiale V, Mourvillier B, Carli P, Varenne O, Mira JP, Wolff M, Cariou A. Early-onset pneumonia after cardiac arrest: characteristics, risk factors and influence on prognosis. *Am J Respir Crit Care Med*. 2011;184:1048–1054. doi: 10.1164/rccm.201102-0331OC
 178. Lybeck A, Friberg H, Aneman A, Hassager C, Horn J, Kjærgaard J, Kuiper M, Nielsen N, Ullén S, Wise MP, Westhall E, Cronberg T; TTM-trial Investigators. Prognostic significance of clinical seizures after cardiac arrest and target temperature management. *Resuscitation*. 2017;114:146–151. doi: 10.1016/j.resuscitation.2017.01.017
 179. Seder DB, Sunde K, Rubertsson S, Mooney M, Ståmmet P, Riker RR, Kern KB, Unger B, Cronberg T, Dziodzio J, et al. Neurologic outcomes and postresuscitation care of patients with myoclonus following cardiac arrest. *Crit Care Med*. 2015;43:965–972. doi: 10.1097/ccm.0000000000000880
 180. Backman S, Westhall E, Dragancea I, Friberg H, Rundgren M, Ullén S, Cronberg T. Electroencephalographic characteristics of status epilepticus after cardiac arrest. *Clin Neurophysiol*. 2017;128:681–688. doi: 10.1016/j.clinph.2017.01.002
 181. Beretta S, Coppo A, Bianchi E, Zanchi C, Carone D, Stabile A, Padovano G, Sulmina E, Grassi A, Bogliun G, Foti G, Ferrarese C, Pesenti A, Beghi E, Avalli L. Neurologic outcome of postanoxic refractory status epilepticus after aggressive treatment. *Neurology*. 2018;91:e2153–e2162. doi: 10.1212/WNL.0000000000006615
 182. Longstreth WT Jr, Fahrenbruch CE, Olsufka M, Walsh TR, Copass MK, Cobb LA. Randomized clinical trial of magnesium, diazepam, or both after out-of-hospital cardiac arrest. *Neurology*. 2002;59:506–514. doi: 10.1212/wnl.59.4.506
 183. Brain Resuscitation Clinical Trial I Study Group. Randomized clinical study of thiopental loading in comatose survivors of cardiac arrest. *New Engl J Med*. 1986;314:397–403. doi: 10.1056/nejm198602133140701
 184. Monsalve F, Rucabado L, Ruano M, Cufat J, Lacueva V, Viñuales A. The neurologic effects of thiopental therapy after cardiac arrest. *Intensive Care Med*. 1987;13:244–248. doi: 10.1007/BF00265112
 185. Thömke F, Weilemann SL. Poor prognosis despite successful treatment of postanoxic generalized myoclonus. *Neurology*. 2010;74:1392–1394. doi: 10.1212/WNL.0b013e3181dad5b9
 186. Koutroumanidis M, Sakellariou D. Low frequency nonevolving generalized periodic epileptiform discharges and the borderland of hypoxic non-convulsive status epilepticus in comatose patients after cardiac arrest. *Epilepsy Behav*. 2015;49:255–262. doi: 10.1016/j.yebeh.2015.04.060
 187. Aicua RI, Rapun I, Novy J, Solari D, Oddo M, Rossetti AO. Early Lance-Adams syndrome after cardiac arrest: prevalence, time to return to awareness, and outcome in a large cohort. *Resuscitation*. 2017;115:169–172. doi: 10.1016/j.resuscitation.2017.03.020
 188. Solanki P, Coppler PJ, Kvaloy JT, Baldwin JM, Callaway CW, Elmer J; Pittsburgh Post-Cardiac Arrest Service. Association of antiepileptic drugs with resolution of epileptiform activity after cardiac arrest. *Resuscitation*. 2019;142:82–90. doi: 10.1016/j.resuscitation.2019.07.007
 189. Kapur J, Elm J, Chamberlain JM, Barsan W, Cloyd J, Lowenstein D, Shinnar S, Conwit R, Meinzer C, Cock H, Fountain N, Connor JT, Silbergleit R; NETT and PECARN Investigators. Randomized Trial of Three Anticonvulsant Medications for Status Epilepticus. *N Engl J Med*. 2019;381:2103–2113. doi: 10.1056/NEJMoa1905795
 190. Elmer J, Rittenberger JC, Faro J, Molyneux BJ, Popescu A, Callaway CW, Baldwin M; Pittsburgh Post-Cardiac Arrest Service. Clinically distinct electroencephalographic phenotypes of early myoclonus after cardiac arrest. *Ann Neurol*. 2016;80:175–184. doi: 10.1002/ana.24697
 191. Crepeau AZ, Fugate JE, Mandrekar J, White RD, Wijdicks EF, Rabinstein AA, Britton JW. Value analysis of continuous EEG in patients during therapeutic hypothermia after cardiac arrest. *Resuscitation*. 2014;85:785–789. doi: 10.1016/j.resuscitation.2014.01.019
 192. Sondag L, Ruijter BJ, Tjepkema-Cloostermans MC, Beishuizen A, Bosch FH, van Til JA, van Putten MJAM, Hofmeijer J. Early EEG for outcome prediction of postanoxic coma: prospective cohort study with cost-minimization analysis. *Crit Care*. 2017;21:111. doi: 10.1186/s13054-017-1693-2
 193. Donnino MW, Andersen LW, Berg KM, Reynolds JC, Nolan JP, Morley PT, Lang E, Cocchi MN, Xanthos T, Callaway CW, Soar J; ILCOR ALS Task Force. Temperature management after cardiac arrest: an advisory statement by the advanced life support task force of the International Liaison Committee on Resuscitation and the American Heart Association Emergency Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. *Resuscitation*. 2016;98:97–104. doi: 10.1016/j.resuscitation.2015.09.396
 194. Donnino MW, Andersen LW, Berg KM, Reynolds JC, Nolan JP, Morley PT, Lang E, Cocchi MN, Xanthos T, Callaway CW, Soar J; ILCOR ALS Task Force. Temperature management after cardiac arrest: an advisory statement by the advanced life support task force of the International Liaison Committee on Resuscitation and the American Heart Association Emergency Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. *Circulation*. 2015;132:2448–2456. doi: 10.1161/CIR.0000000000000313
 195. Lascarrrou JB, Merdji H, Le Gouge A, Colin G, Grillet G, Girardie P, Coupez E, Dequin PF, Cariou A, Boulain T, Brule N, Frat JP, Asfar P, Pichon N, Landais M, Plantefève G, Quenot JP, Chakarian JC, Sirodot M, Legriel S, Lethuelle J, Thevenin D, Desachy A, Delahaye A, Botoc V, Vimeux S, Martino F, Giraudeau B, Reignier J; CRICS-TRIGGERSEP Group. Targeted Temperature Management for Cardiac Arrest with Nonshockable Rhythm. *N Engl J Med*. 2019;381:2327–2337. doi: 10.1056/NEJMoa1906661
 196. Sandroni C, D'Arrigo S, Cacciola S, Hoedemaekers CWE, Kamps MJA, Oddo M, Taccone FS, Di Rocco A, Meijer FJA, Westhall E, et al. Prediction of poor neurological outcome in comatose survivors of cardiac arrest: a systematic review [published online September 11, 2020]. *Intensive Care Med*. doi: 10.1007/s00134-020-06198-w
 197. Steinberg A, Callaway CW, Arnold RM, Cronberg T, Naito H, Dadon K, Chae MK, Elmer J. Prognostication after cardiac arrest: results of an international, multi-professional survey. *Resuscitation*. 2019;138:190–197. doi: 10.1016/j.resuscitation.2019.03.016
 198. Ryoo SM, Jeon SB, Sohn CH, Ahn S, Han C, Lee BK, Lee DH, Kim SH, Donnino MW, Kim WY; Korean Hypothermia Network Investigators. Predicting outcome with diffusion-weighted imaging in cardiac arrest patients receiving hypothermia therapy: multicenter retrospective cohort study. *Crit Care Med*. 2015;43:2370–2377. doi: 10.1097/CCM.0000000000001263
 199. Scarpino M, Lolli F, Lanzo G, Carrai R, Spalletti M, Valzania F, Lombardi M, Audenino D, Celani MG, Marrelli A, Contardi S, Peris A, Amantini A, Sandroni C, Grippo A; ProNeCASTudy Group. Neurophysiological and neuroradiological test for early poor outcome (Cerebral Performance Categories 3-5) prediction after cardiac arrest: Prospective multicentre prognostication data. *Data Brief*. 2019;27:104755. doi: 10.1016/j.dib.2019.104755
 200. Kongpoltprom N, Cholkraisuwat J. Neurological prognostications for the therapeutic hypothermia among comatose survivors of cardiac arrest. *Indian J Crit Care Med*. 2018;22:509–518. doi: 10.4103/ijccm.IJCCM_500_17
 201. Roger C, Palmier L, Louart B, Molinari N, Claret PG, de la Coussaye JE, Lefrant JY, Muller L. Neuron specific enolase and Glasgow motor score remain useful tools for assessing neurological prognosis after out-of-hospital

- cardiac arrest treated with therapeutic hypothermia. *Anaesth Crit Care Pain Med*. 2015;34:231–237. doi: 10.1016/j.accpm.2015.05.004
202. Zhou SE, Maciel CB, Ormseth CH, Beekman R, Gilmore EJ, Greer DM. Distinct predictive values of current neuroprognostic guidelines in post-cardiac arrest patients. *Resuscitation*. 2019;139:343–350. doi: 10.1016/j.resuscitation.2019.03.035
 203. Lee KS, Lee SE, Choi JY, Cho YR, Chae MK, Park EJ, Choi MH, Hong JM. Useful computed tomography score for estimation of early neurologic outcome in post-cardiac arrest patients with therapeutic hypothermia. *Circ J*. 2017;81:1628–1635. doi: 10.1253/circj.CJ-16-1327
 204. Matthews EA, Magid-Bernstein J, Sobczak E, Velazquez A, Falo CM, Park S, Claassen J, Agarwal S. Prognostic Value of the Neurological Examination in Cardiac Arrest Patients After Therapeutic Hypothermia. *Neurohospitalist*. 2018;8:66–73. doi: 10.1177/1941874417733217
 205. Choi SP, Park KN, Wee JH, Park JH, Youn CS, Kim HJ, Oh SH, Oh JS, Kim SH, Oh JS. Can somatosensory and visual evoked potentials predict neurological outcome during targeted temperature management in post cardiac arrest patients? *Resuscitation*. 2017;119:70–75. doi: 10.1016/j.resuscitation.2017.06.022
 206. Chung-Esaki HM, Mui G, Mlynash M, Eynogrn I, Catabay K, Hirsch KG. The neuron specific enolase (NSE) ratio offers benefits over absolute value thresholds in post-cardiac arrest coma prognosis. *J Clin Neurosci*. 2018;57:99–104. doi: 10.1016/j.jocn.2018.08.020
 207. Javaudin F, Leclere B, Segard J, Le Bastard Q, Pes P, Penverne Y, Le Conte P, Jenvrin J, Hubert H, Escutnaire J, Bataud E, Montassier E, Gr-RéAC. Prognostic performance of early absence of pupillary light reaction after recovery of out of hospital cardiac arrest. *Resuscitation*. 2018;127:8–13. doi: 10.1016/j.resuscitation.2018.03.020
 208. Dhakal LP, Sen A, Stanko CM, Rawal B, Heckman MG, Hoyne JB, Dimberg EL, Freeman ML, Ng LK, Rabinstein AA, Freeman WD. Early absent pupillary light reflexes after cardiac arrest in patients treated with therapeutic hypothermia. *Ther Hypothermia Temp Manag*. 2016;6:116–121. doi: 10.1089/ther.2015.0035
 209. Oddo M, Sandroni C, Citerio G, Miroz JP, Horn J, Rundgren M, Cariou A, Payen JF, Storm C, Stamment P, Taccone FS. Quantitative versus standard pupillary light reflex for early prognostication in comatose cardiac arrest patients: an international prospective multicenter double-blinded study. *Intensive Care Med*. 2018;44:2102–2111. doi: 10.1007/s00134-018-5448-6
 210. Fatuzzo D, Beuchat I, Alvarez V, Novy J, Oddo M, Rossetti AO. Does continuous EEG influence prognosis in patients after cardiac arrest? *Resuscitation*. 2018;132:29–32. doi: 10.1016/j.resuscitation.2018.08.023
 211. Draganca I, Horn J, Kuiper M, Friberg H, Ullén S, Wetterslev J, Cranshaw J, Hassager C, Nielsen N, Cronberg T; TTM Trial Investigators. Neurological prognostication after cardiac arrest and targeted temperature management 33°C versus 36°C: Results from a randomised controlled clinical trial. *Resuscitation*. 2015;93:164–170. doi: 10.1016/j.resuscitation.2015.04.013
 212. Hofmeijer J, Beernink TM, Bosch FH, Beishuizen A, Tjepkema-Cloostermans MC, van Putten MJ. Early EEG contributes to multimodal outcome prediction of postanoxic coma. *Neurology*. 2015;85:137–143. doi: 10.1212/WNL.0000000000001742
 213. Greer DM, Yang J, Scripko PD, Sims JR, Cash S, Wu O, Hafner JP, Schoenfeld DA, Furie KL. Clinical examination for prognostication in comatose cardiac arrest patients. *Resuscitation*. 2013;84:1546–1551. doi: 10.1016/j.resuscitation.2013.07.028
 214. Kim JH, Kim MJ, You JS, Lee HS, Park YS, Park I, Chung SP. Multimodal approach for neurologic prognostication of out-of-hospital cardiac arrest patients undergoing targeted temperature management. *Resuscitation*. 2019;134:33–40. doi: 10.1016/j.resuscitation.2018.11.007
 215. Solari D, Rossetti AO, Carteron L, Miroz JP, Novy J, Eckert P, Oddo M. Early prediction of coma recovery after cardiac arrest with blinded pupillometry. *Ann Neurol*. 2017;81:804–810. doi: 10.1002/ana.24943
 216. Heimbürger D, Durand M, Gaide-Chevronnay L, Dessertaine G, Moury PH, Bouzat P, Albaladejo P, Payen JF. Quantitative pupillometry and transcranial Doppler measurements in patients treated with hypothermia after cardiac arrest. *Resuscitation*. 2016;103:88–93. doi: 10.1016/j.resuscitation.2016.02.026
 217. Riker RR, Sawyer ME, Fischman VG, May T, Lord C, Eldridge A, Seder DB. Neurological Pupil Index and Pupillary Light Reflex by Pupillometry Predict Outcome Early After Cardiac Arrest. *Neurocrit Care*. 2020;32:152–161. doi: 10.1007/s12028-019-00717-4
 218. Obling L, Hassager C, Illum C, Grand J, Wiberg S, Lindholm MG, Winther-Jensen M, Kondziella D, Kjaergaard J. Prognostic value of automated pupillometry: an unselected cohort from a cardiac intensive care unit. *Eur Heart J Acute Cardiovasc Care*. 2019;20:48872619842004. doi: 10.1177/2048872619842004
 219. Sivaraju A, Gilmore EJ, Wira CR, Stevens A, Rampal N, Moeller JJ, Greer DM, Hirsch LJ, Gaspard N. Prognostication of post-cardiac arrest coma: early clinical and electroencephalographic predictors of outcome. *Intensive Care Med*. 2015;41:1264–1272. doi: 10.1007/s00134-015-3834-x
 220. Sadaka F, Doerr D, Hindia J, Lee KP, Logan W. Continuous electroencephalogram in comatose postcardiac arrest syndrome patients treated with therapeutic hypothermia: outcome prediction study. *J Intensive Care Med*. 2015;30:292–296. doi: 10.1177/0885066613517214
 221. Maia B, Roque R, Amaral-Silva A, Lourenço S, Bento L, Alcântara J. Predicting outcome after cardiopulmonary arrest in therapeutic hypothermia patients: clinical, electrophysiological and imaging prognosticators. *Acta Med Port*. 2013;26:93–97.
 222. Reynolds AS, Rohaut B, Holmes MG, Robinson D, Roth W, Velazquez A, Couch CK, Presciutti A, Brodie D, Moitra VK, Rabbani LE, Agarwal S, Park S, Roh DJ, Claassen J. Early myoclonus following anoxic brain injury. *Neurolog Clin Pract*. 2018;8:249–256. doi: 10.1212/CPJ.0000000000000466
 223. Ruknudeen MI, Ramadoss R, Rajajee V, Grzeskowiak LE, Rajaopalan RE. Early clinical prediction of neurological outcome following out of hospital cardiac arrest managed with therapeutic hypothermia. *Indian J Crit Care Med*. 2015;19:304–310. doi: 10.4103/0972-5229.158256
 224. Maciel CB, Morawo AO, Tsao CY, Youn TS, Labar DR, Rubens EO, Greer DM. SSEP in therapeutic hypothermia era. *J Clin Neurophysiol*. 2017;34:469–475. doi: 10.1097/WNP.0000000000000392
 225. Leão RN, Ávila P, Cavaco R, Germano N, Bento L. Therapeutic hypothermia after cardiac arrest: outcome predictors. *Rev Bras Ter Intensiva*. 2015;27:322–332. doi: 10.5935/0103-507X.20150056
 226. Ruijter BJ, Tjepkema-Cloostermans MC, Tromp SC, van den Bergh WM, Foudraine NA, Kornips FHM, Drost G, Scholten E, Bosch FH, Beishuizen A, van Putten MJAM, Hofmeijer J. Early electroencephalography for outcome prediction of postanoxic coma: A prospective cohort study. *Ann Neurol*. 2019;86:203–214. doi: 10.1002/ana.25518
 227. Kim SW, Oh JS, Park J, Jeong HH, Kim JH, Wee JH, Oh SH, Choi SP, Park KN; Cerebral Resuscitation and Outcome evaluation Within catholic Network (CROWN) Investigators. Short-latency positive peak following N20 somatosensory evoked potential is superior to N20 in predicting neurologic outcome after out-of-hospital cardiac arrest. *Crit Care Med*. 2018;46:e545–e551. doi: 10.1097/CCM.00000000000003083
 228. Scarpino M, Carrai R, Lolli F, Lanzo G, Spalletti M, Valzania F, Lombardi M, Audenino D, Contardi S, Celani MG, Marrelli A, Mecarelli O, Minardi C, Minicucci F, Politini L, Vitelli E, Peris A, Amantini A, Sandroni C, Grippo A; ProNeCA study group. Neurophysiology for predicting good and poor neurological outcome at 12 and 72 h after cardiac arrest: The ProNeCA multicentre prospective study. *Resuscitation*. 2020;147:95–103. doi: 10.1016/j.resuscitation.2019.11.014
 229. Grippo A, Carrai R, Scarpino M, Spalletti M, Lanzo G, Cossu C, Peris A, Valente S, Amantini A. Neurophysiological prediction of neurological good and poor outcome in post-anoxic coma. *Acta Neurol Scand*. 2017;135:641–648. doi: 10.1111/ane.12659
 230. De Santis P, Lamanna I, Mavroudikis N, Legros B, Vincent JL, Creteur J, Taccone FS. The potential role of auditory evoked potentials to assess prognosis in comatose survivors from cardiac arrest. *Resuscitation*. 2017;120:119–124. doi: 10.1016/j.resuscitation.2017.09.013
 231. Westhäll E, Rossetti AO, van Rootselaar AF, Wesenberg Kjaer T, Horn J, Ullén S, Friberg H, Nielsen N, Rosén I, Åneman A, Erlinge D, Gasche Y, Hassager C, Hovdenes J, Kjaergaard J, Kuiper M, Pellis T, Stamment P, Wanscher M, Wetterslev J, Wise MP, Cronberg T; TTM-trial investigators. Standardized EEG interpretation accurately predicts prognosis after cardiac arrest. *Neurology*. 2016;86:1482–1490. doi: 10.1212/WNL.0000000000002462
 232. Admiraal MM, van Rootselaar AF, Hofmeijer J, Hoedemakers CWE, van Kaam CR, Keijzer HM, van Putten MJAM, Schultz MJ, Horn J. Electroencephalographic reactivity as predictor of neurological outcome in postanoxic coma: A multicenter prospective cohort study. *Ann Neurol*. 2019;86:17–27. doi: 10.1002/ana.25507
 233. Alvarez V, Reinsberger C, Scirica B, O'Brien MH, Avery KR, Henderson G, Lee JW. Continuous electrodermal activity as a potential novel neurophysiological biomarker of prognosis after cardiac arrest—A pilot study. *Resuscitation*. 2015;93:128–135. doi: 10.1016/j.resuscitation.2015.06.006
 234. Duez CHV, Johnsen B, Ebbesen MQ, Kvaløy MB, Grejs AM, Jeppesen AN, Søreide E, Nielsen JF, Kirkegaard H. Post resuscitation prognostication by EEG in 24 vs 48 h of targeted temperature management. *Resuscitation*. 2019;135:145–152. doi: 10.1016/j.resuscitation.2018.10.035

235. Liu G, Su Y, Liu Y, Jiang M, Zhang Y, Zhang Y, Gao D. Predicting outcome in comatose patients: the role of EEG reactivity to quantifiable electrical stimuli. *Evid Based Complement Alternat Med*. 2016;2016:8273716. doi: 10.1155/2016/8273716
236. Amorim E, Rittenberger JC, Zheng JJ, Westover MB, Baldwin ME, Callaway CW, Popescu A; Post Cardiac Arrest Service. Continuous EEG monitoring enhances multimodal outcome prediction in hypoxic-ischemic brain injury. *Resuscitation*. 2016;109:121–126. doi: 10.1016/j.resuscitation.2016.08.012
237. Benarous L, Gavaret M, Soda Diop M, Tobarías J, de Ghaisne de Bourmont S, Allez C, Bouzana F, Gainnier M, Trebuchon A. Sources of interrater variability and prognostic value of standardized EEG features in post-anoxic coma after resuscitated cardiac arrest. *Clin Neurophysiol Pract*. 2019;4:20–26. doi: 10.1016/j.cnp.2018.12.001
238. Lamartine Monteiro M, Taccone FS, Depondt C, Lamanna I, Gaspard N, Ligot N, Mavroudikis N, Naeije G, Vincent JL, Legros B. The prognostic value of 48-h continuous EEG during therapeutic hypothermia after cardiac arrest. *Neurocrit Care*. 2016;24:153–162. doi: 10.1007/s12028-015-0215-9
239. Rossetti AO, Tovar Quiroga DF, Juan E, Novy J, White RD, Ben-Hamouda N, Britton JW, Oddo M, Rabinstein AA. Electroencephalography predicts poor and good outcomes after cardiac arrest: a two-center study. *Crit Care Med*. 2017;45:e674–e682. doi: 10.1097/CCM.0000000000002337
240. Backman S, Cronberg T, Friberg H, Ullén S, Horn J, Kjaergaard J, Hassager C, Wanscher M, Nielsen N, Westhall E. Highly malignant routine EEG predicts poor prognosis after cardiac arrest in the Target Temperature Management trial. *Resuscitation*. 2018;131:24–28. doi: 10.1016/j.resuscitation.2018.07.024
241. Beretta S, Coppo A, Bianchi E, Zanchi C, Carone D, Stabile A, Padovano G, Sulmina E, Grassi A, Bogliun G, Foti G, Ferrarese C, Pesenti A, Beghi E, Avalli L. Neurological outcome of postanoxic refractory status epilepticus after aggressive treatment. *Epilepsy Behav*. 2019;101(Pt B):106374. doi: 10.1016/j.yebeh.2019.06.018
242. Oh SH, Park KN, Shon YM, Kim YM, Kim HJ, Youn CS, Kim SH, Choi SP, Kim SC. Continuous Amplitude-Integrated Electroencephalographic Monitoring Is a Useful Prognostic Tool for Hypothermia-Treated Cardiac Arrest Patients. *Circulation*. 2015;132:1094–1103. doi: 10.1161/CIRCULATIONAHA.115.015754
243. Dragancea I, Backman S, Westhall E, Rundgren M, Friberg H, Cronberg T. Outcome following postanoxic status epilepticus in patients with targeted temperature management after cardiac arrest. *Epilepsy Behav*. 2015;49:173–177. doi: 10.1016/j.yebeh.2015.04.043
244. Hirsch LJ, LaRoche SM, Gaspard N, Gerard E, Svoronos A, Herman ST, Mani R, Arif H, Jette N, Minazad Y, et al. American Clinical Neurophysiology Society's Standardized Critical Care EEG Terminology: 2012 version. *J Clin Neurophysiol*. 2013;30:1–27. doi: 10.1097/WNP.0b013e3182784729
245. Admiraal MM, van Rootselaar AF, Horn J. International consensus on EEG reactivity testing after cardiac arrest: towards standardization. *Resuscitation*. 2018;131:36–41. doi: 10.1016/j.resuscitation.2018.07.025
246. Zellner T, Gärtner R, Schopohl J, Angstwurm M. NSE and S-100B are not sufficiently predictive of neurologic outcome after therapeutic hypothermia for cardiac arrest. *Resuscitation*. 2013;84:1382–1386. doi: 10.1016/j.resuscitation.2013.03.021
247. Lee BK, Jeung KW, Lee HY, Jung YH, Lee DH. Combining brain computed tomography and serum neuron specific enolase improves the prognostic performance compared to either alone in comatose cardiac arrest survivors treated with therapeutic hypothermia. *Resuscitation*. 2013;84:1387–1392. doi: 10.1016/j.resuscitation.2013.05.026
248. Vondrakova D, Kruger A, Janotka M, Malek F, Dudkova V, Neuzil P, Ostadal P. Association of neuron-specific enolase values with outcomes in cardiac arrest survivors is dependent on the time of sample collection. *Crit Care*. 2017;21:172. doi: 10.1186/s13054-017-1766-2
249. Stammet P, Collignon O, Hassager C, Wise MP, Hovdenes J, Aneman A, Horn J, Devaux Y, Erlinge D, Kjaergaard J, Gasche Y, Wanscher M, Cronberg T, Friberg H, Wetterslev J, Pellis T, Kuiper M, Gilson G, Nielsen N; TTM-Trial Investigators. Neuron-specific enolase as a predictor of death or poor neurological outcome after out-of-hospital cardiac arrest and targeted temperature management at 33°C and 36°C. *J Am Coll Cardiol*. 2015;65:2104–2114. doi: 10.1016/j.jacc.2015.03.538
250. Tsetsou S, Novy J, Pfeiffer C, Oddo M, Rossetti AO. Multimodal outcome prognostication after cardiac arrest and targeted temperature management: analysis at 36 °C. *Neurocrit Care*. 2018;28:104–109. doi: 10.1007/s12028-017-0393-8
251. Duez CHV, Grejs AM, Jeppesen AN, Schrøder AD, Sørreide E, Nielsen JF, Kirkegaard H. Neuron-specific enolase and S-100b in prolonged targeted temperature management after cardiac arrest: A randomised study. *Resuscitation*. 2018;122:79–86. doi: 10.1016/j.resuscitation.2017.11.052
252. Helwig K, Seeger F, Hölschermann H, Lischke V, Gerriets T, Niessner M, Foerch C. Elevated serum glial fibrillary acidic protein (GFAP) is associated with poor functional outcome after cardiopulmonary resuscitation. *Neurocrit Care*. 2017;27:68–74. doi: 10.1007/s12028-016-0371-6
253. Stammet P, Dankiewicz J, Nielsen N, Fays F, Collignon O, Hassager C, Wanscher M, Undèn J, Wetterslev J, Pellis T, Aneman A, Hovdenes J, Wise MP, Gilson G, Erlinge D, Horn J, Cronberg T, Kuiper M, Kjaergaard J, Gasche Y, Devaux Y, Friberg H; Target temperature management after out-of-hospital cardiac arrest (TTM) trial investigators. Protein S100 as outcome predictor after out-of-hospital cardiac arrest and targeted temperature management at 33 °C and 36 °C. *Crit Care*. 2017;21:153. doi: 10.1186/s13054-017-1729-7
254. Jang JH, Park WB, Lim YS, Choi JY, Cho JS, Woo JH, Choi WS, Yang HJ, Hyun SY. Combination of S100B and procalcitonin improves prognostic performance compared to either alone in patients with cardiac arrest: a prospective observational study. *Medicine (Baltimore)*. 2019;98:e14496. doi: 10.1097/MD.00000000000014496
255. Mattsson N, Zetterberg H, Nielsen N, Blennow K, Dankiewicz J, Friberg H, Lilja G, Insel PS, Rylander C, Stammet P, Aneman A, Hassager C, Kjaergaard J, Kuiper M, Pellis T, Wetterslev J, Wise M, Cronberg T. Serum tau and neurological outcome in cardiac arrest. *Ann Neurol*. 2017;82:665–675. doi: 10.1002/ana.25067
256. Moseby-Knappe M, Mattsson N, Nielsen N, Zetterberg H, Blennow K, Dankiewicz J, Dragancea I, Friberg H, Lilja G, Insel PS, Rylander C, Westhall E, Kjaergaard J, Wise MP, Hassager C, Kuiper MA, Stammet P, Wanscher MCJ, Wetterslev J, Erlinge D, Horn J, Pellis T, Cronberg T. Serum neurofilament light chain for prognosis of outcome after cardiac arrest. *JAMA Neurol*. 2019;76:64–71. doi: 10.1001/jamaneurol.2018.3223
257. Rana OR, Schröder JW, Baukloh JK, Saygili E, Mischke K, Schiefer J, Weis J, Marx N, Rassaf T, Kelm M, Shin DI, Meyer C, Saygili E. Neurofilament light chain as an early and sensitive predictor of long-term neurological outcome in patients after cardiac arrest. *Int J Cardiol*. 2013;168:1322–1327. doi: 10.1016/j.ijcard.2012.12.016
258. Wang GN, Chen XF, Lv JR, Sun NN, Xu XQ, Zhang JS. The prognostic value of gray-white matter ratio on brain computed tomography in adult comatose cardiac arrest survivors. *J Chin Med Assoc*. 2018;81:599–604. doi: 10.1016/j.jcma.2018.03.003
259. Youn CS, Callaway CW, Rittenberger JC; Post Cardiac Arrest Service. Combination of initial neurologic examination, quantitative brain imaging and electroencephalography to predict outcome after cardiac arrest. *Resuscitation*. 2017;110:120–125. doi: 10.1016/j.resuscitation.2016.10.024
260. Jeon CH, Park JS, Lee JH, Kim H, Kim SC, Park KH, Yi KS, Kim SM, Youn CS, Kim YM, Lee BK. Comparison of brain computed tomography and diffusion-weighted magnetic resonance imaging to predict early neurologic outcome before target temperature management comatose cardiac arrest survivors. *Resuscitation*. 2017;118:21–26. doi: 10.1016/j.resuscitation.2017.06.021
261. Kim SH, Choi SP, Park KN, Youn CS, Oh SH, Choi SM. Early brain computed tomography findings are associated with outcome in patients treated with therapeutic hypothermia after out-of-hospital cardiac arrest. *Scand J Trauma Resusc Emerg Med*. 2013;21:57. doi: https://doi.org/10.1186/1757-7241-21-57
262. Kim Y, Ho LJ, Kun HC, Won CK, Hoon YJ, Ju KM, Weon KY, Yul LK, Joo KJ, Youn HS. Feasibility of optic nerve sheath diameter measured on initial brain computed tomography as an early neurologic outcome predictor after cardiac arrest. *Academic Emergency Medicine* 2014;21:1121–1128.
263. Kim SJ, Jung JS, Park JH, Park JS, Hong YS, Lee SW. An optimal transition time to extracorporeal cardiopulmonary resuscitation for predicting good neurological outcome in patients with out-of-hospital cardiac arrest: a propensity-matched study. *Crit Care*. 2014;18:535. doi: 10.1186/s13054-014-0535-8
264. Scarpino M, Lanzò G, Lolli F, Carrai R, Moretti M, Spalletti M, Cozzolino M, Peris A, Amantini A, Grippo A. Neurophysiological and neuroradiological multimodal approach for early poor outcome prediction after cardiac arrest. *Resuscitation*. 2018;129:114–120. doi: 10.1016/j.resuscitation.2018.04.016
265. Lee DH, Lee BK, Jeung KW, Jung YH, Cho YS, Cho IS, Youn CS, Kim JW, Park JS, Min YI. Relationship between ventricular characteristics on brain computed tomography and 6-month neurological outcome in cardiac arrest survivors who underwent targeted temperature management. *Resuscitation*. 2018;129:37–42. doi: 10.1016/j.resuscitation.2018.06.008
266. Lee BK, Jeung KW, Song KH, Jung YH, Choi WJ, Kim SH, Youn CS, Cho IS, Lee DH; Korean Hypothermia Network Investigators. Prognostic

- values of gray matter to white matter ratios on early brain computed tomography in adult comatose patients after out-of-hospital cardiac arrest of cardiac etiology. *Resuscitation*. 2015;96:46–52. doi: 10.1016/j.resuscitation.2015.07.027
267. Lee BK, Kim WY, Shin J, Oh JS, Wee JH, Cha KC, Park Y, Choi JH, Jeung KW; Korean Hypothermia Network Investigators. Prognostic value of gray matter to white matter ratio in hypoxic and non-hypoxic cardiac arrest with non-cardiac etiology. *Am J Emerg Med*. 2016;34:1583–1588. doi: 10.1016/j.ajem.2016.05.063
268. Greer DM, Scripko PD, Wu O, Edlow BL, Bartscher J, Sims JR, Camargo EE, Singhal AB, Furie KL. Hippocampal magnetic resonance imaging abnormalities in cardiac arrest are associated with poor outcome. *J Stroke Cerebrovasc Dis*. 2013;22:899–905. doi: 10.1016/j.jstrokecerebrovasdis.2012.08.006
269. Jang J, Oh SH, Nam Y, Lee K, Choi HS, Jung SL, Ahn KJ, Park KN, Kim BS. Prognostic value of phase information of 2D T2*-weighted gradient echo brain imaging in cardiac arrest survivors: A preliminary study. *Resuscitation*. 2019;140:142–149. doi: 10.1016/j.resuscitation.2019.05.026
- 269a. Kim J, Kim K, Hong S, Kwon B, Yun ID, Choi BS, Jung C, Lee JH, Jo YH, Kim T, et al. Low apparent diffusion coefficient cluster-based analysis of diffusion-weighted MRI for prognostication of out-of-hospital cardiac arrest survivors. *Resuscitation*. 2013;84:1393–1399. doi: 10.1016/j.resuscitation.2013.04.01
270. Moon HK, Jang J, Park KN, Kim SH, Lee BK, Oh SH, Jeung KW, Choi SP, Cho IS, Youn CS. Quantitative analysis of relative volume of low apparent diffusion coefficient value can predict neurologic outcome after cardiac arrest. *Resuscitation*. 2018;126:36–42. doi: 10.1016/j.resuscitation.2018.02.020
271. Nikolaou NI, Welsford M, Beygui F, Bossaert L, Ghaemmaghami C, Nonogi H, O'Connor RE, Pichel DR, Scott T, Walters DL, et al; on behalf of the Acute Coronary Syndrome Chapter Collaborators. Part 5: acute coronary syndromes: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2015;95:e121–e146. doi: 10.1016/j.resuscitation.2015.07.043
272. Welsford M, Nikolaou NI, Beygui F, Bossaert L, Ghaemmaghami C, Nonogi H, O'Connor RE, Pichel DR, Scott T, Walters DL, et al; on behalf of the Acute Coronary Syndrome Chapter Collaborators. Part 5: acute coronary syndromes: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2015;132(suppl 1):S146–S176. doi: 10.1161/CIR.0000000000000274